

Publications Désirée van der Heijde

1. Thesis

1991

Disease activity and outcome in rheumatoid arthritis: a methodological study.

Katholieke Universiteit Nijmegen 8 march 1991

2. Scientific international papers in relation to imaging

2018

van der Heijde D, Baraliakos X, Hermann K-GA, Landewé RBM, Machado PM, Maksymowych WP, Davies OR, de Peyrecave N, Hoepken B, Bauer L, Nurminen T, Braun J. Limited radiographic progression and sustained reductions in MRI inflammation in patients with axial spondyloarthritis: 4-year imaging outcomes from the RAPID-axSpA phase III randomised trial.

Ann Rheum Dis 2018 Online first

de Bruin F, de Koning A, van den Berg R, Baraliakos X, Braun J, Ramiro S, van Gaalen FA, Reijnierse M, van der Heijde D. Development of the CT Syndesmophyte Score (CTSS) in patients with ankylosing spondylitis: data from the SIAS cohort.

Ann Rheum Dis 2018;77:371–7

de Koning A, de Bruin F, van den Berg R, Ramiro S, Baraliakos X, Braun J, van Gaalen FA, Reijnierse M, van der Heijde D. Low-dose CT detects more progression of bone formation in comparison to conventional radiography in patients with ankylosing spondylitis: results from the SIAS cohort.

Ann Rheum Dis 2018;77:293–99

Molnar C, Scherer A, Baraliakos X, de Hooge M, Micheroli R, Exer P, Kissling RO, Tamborrini G, Wildi LM, Nissen MJ, Zufferey P, Bernhard J, Weber U, Landewé RBM, van der Heijde D, Ciurea A. TNF blockers inhibit spinal radiographic progression in ankylosing spondylitis by reducing disease activity: results from the Swiss Clinical Quality Management cohort.

Ann Rheum Dis 2018;77:63-9

2017

van der Heijde D, Gladman DD, Kishimoto M, Okada M, Rathmann SS, Moriarty SR, Shuler CL, Carlier H, Benichou O, Mease PJ. Efficacy and Safety of Ixekizumab in Patients with Active Psoriatic Arthritis: 52-week Results from a Phase III Study (SPIRIT-P1).

J Rheumatol. 2017 Dec 15. pii: jrheum.170429. doi: 10.3899/jrheum.170429. [Epub ahead of print]

Charles-Schoeman C, van der Heijde D, Burmester GR, Nash P, Zerbini CAF, Connell CA, Fan H, Kwok K, Bananis E, Fleischmann R. Effect of Glucocorticoids on the Clinical and Radiographic Efficacy of Tofacitinib in Patients with Rheumatoid Arthritis: A Posthoc Analysis of Data from 6 Phase III Studies.

J Rheumatology 2017, jrheum.170486; DOI: <https://doi.org/10.3899/jrheum.170486>

de Bruin F, ter Horst S, Bloem JL, van den Berg R, de Hooze M, van Gaalen F, Dagfinrud H, van Oosterhout M, Landewé R, van der Heijde D, Reijnen M. Prevalence and clinical significance of lumbosacral transitional vertebra (LSTV) in a young back pain population with suspected axial spondyloarthritis: results of the SPondyloArthritis Caught Early (SPACE) cohort. Skeletal Radiol 2017;46:633–9

de Hooze M, Pialat JB, Reijnen M, van der Heijde D, Claudepierre P, Saraux A, Dougados M, Feydy A. Assessment of typical SpA lesions on MRI of the spine: do local readers and central readers agree in the DESIR-cohort at baseline? Clin Rheumatol. 2017 Jul;36(7):1551-1559

Braun J, Baraliakos X, Hermann KG, Landewé R, Machado PM, Maksymowych WP, Davies O, Hoepken B, Nurminen T, Stach C, van der Heijde D. Effect of certolizumab pegol over 96 weeks of treatment on inflammation of the spine and sacroiliac joints, as measured by MRI, and the association between clinical and MRI outcomes in patients with axial spondyloarthritis.

RMD Open 2017;3:e000430

Mease P, Gottlieb AB, van der Heijde D, FitzGerald O, Johnsen A, Nys M, Banerjee S, Gladman DD. Efficacy and safety of abatacept, a T-cell modulator, in a randomised, double-blind, placebo-controlled, phase III study in psoriatic arthritis. Ann Rheum Dis 2017;76:1550-8

de Hooze M, de Bruin F, de Beer L, Bakker P, van den Berg R, Ramiro S, van Gaalen F, Fagerli K, Landewé R, van Oosterhout M, Ramonda R, Huizinga T, Bloem H, Reijnen M, van der Heijde D. Is site of back pain related to location of MRI lesion in patients with chronic back pain included in the SPACE-cohort?

Arthritis Care Res 2017;69:717-23

Haugen IK, Slatkowsky-Christensen B, Faraj K, van der Heijde D, Kvien TK. The comparison of magnetic resonance imaging and radiographs to assess structural progression over 5 years in hand osteoarthritis.

Rheumatology 2017;56:371-7

Aizenberg E, van den Berg R, Ez-Zaitouni Z, van der Heijde D, Reijnen M, Dzyubachyk O, Lelieveldt BPF. Computer-aided evaluation of inflammatory changes over time on MRI of the spine in patients with suspected axial spondyloarthritis: a feasibility study.

BMC Medical Imaging 2017;17:55

Mease P, Hall S, FitzGerald O, van der Heijde D, Merola JF, Avila-Zapata F, Cieślak D, Graham D, Wang C, Menon S, Hendrikx T, Kanik KS. Tofacitinib or Adalimumab versus Placebo for Psoriatic Arthritis.

N Engl J Med 2017; 377:1537-50

Brinkmann GH, Norli ES, Bøyesen P, [van der Heijde D](#), Grøvlø L, Haugen AJ, Nygaard H, Bjørneboe O, Thunem C, Kvien TK, Mjaavatten MD, Lie E. Role of erosions typical of rheumatoid arthritis in the 2010 ACR/EULAR rheumatoid arthritis classification criteria: results from a very early arthritis cohort. *Ann Rheum Dis* 2017;76:1911-4

Dougados M, Sepriano A, Molto A, van Lunteren M, Ramiro S, de Hooge M, van den Berg R, Navarro Compan V, Demattei C, Landewé R, [van der Heijde D](#). Sacroiliac radiographic progression in recent onset axial spondyloarthritis: the 5-year data of the DESIR cohort. *Ann Rheum Dis* 2017;76:1823-8

Ez-Zaitouni Z, Bakker PAC, van Lunteren M, de Hooge M, van den Berg R, Reijniere M, Fagerli KM, Landewé RBM, Ramonda R, Jacobsson LTH, Saraux A, Lenczner G, Feydy A, Pialat JB, Thévenin F, van Gaalen FA, [van der Heijde D](#). The yield of a positive MRI of the spine as imaging criterion in the ASAS classification criteria for axial spondyloarthritis: results from the SPACE and DESIR cohorts. *Ann Rheum Dis* 2017; 76:1731-6

Atsumi T, Tanaka Y, Yamamoto K, Takeuchi T, Yamanaka H, Ishiguro N, Eguchi K, Watanabe A, Origasa H, Yasuda S, Yamanishi Y, Kita Y, Matsubara T, Iwamoto M, Shoji T, Togo O, Okada T, [van der Heijde D](#), Miyasaka N, Koike T. Clinical benefit of 1-year certolizumab pegol (CZP) add-on therapy to methotrexate treatment in patients with early rheumatoid arthritis was observed following CZP discontinuation: 2-year results of the C-OPERA study, a phase III randomised trial. *Ann Rheum Dis* 2017;76:1348-56

[van der Heijde D](#), Deodhar A, Wei JC, Drescher E, Fleishaker D, Hendriks T, Li D, Menon S, Kanik KS. Tofacitinib in patients with ankylosing spondylitis: a phase II, 16-week, randomised, placebo-controlled, dose-ranging study. *Ann Rheum Dis* 2017;76:1340-7

Taylor PC, Keystone EC, [van der Heijde D](#), Weinblatt ME, del Carmen Morales L, Reyes Gonzaga J, Yakushin S, Ishii T, Emoto K, Beattie S, Arora V, Gaich C, Rooney T, Schlichting D, Macias WL, de Bono S, Tanaka Y. Baricitinib versus Placebo or Adalimumab in Rheumatoid Arthritis. *N Engl J Med* 2017; 376:652-62

Bakker PAC, van den Berg R, Lenczner G, Thévenin F, Reijniere M, Claudepierre P, Wendling D, Dougados M, [van der Heijde D](#). Can we use structural lesions seen on MRI of the sacroiliac joints reliably for the classification of patients according to the ASAS axial spondyloarthritis criteria? Data from the DESIR cohort. *Ann Rheum Dis* 2017; 76:392-8

Emery P, Bingham CO III, Burmester GR, Bykerk VP, Furst DE, Mariette X, [van der Heijde D](#), van Vollenhoven R, Arendt C, Moutian I, Purcaru O, Tatla D, VanLunen B, Weinblatt ME. Certolizumab pegol in combination with dose-optimised methotrexate in DMARD-naïve patients with early, active rheumatoid arthritis with poor prognostic factors: 1-year results from C-EARLY, a randomised, double-blind, placebo-controlled phase III study. *Ann Rheum Dis* 2017;76:96-104

Mease PJ, [van der Heijde D](#), Ritchlin CT, Okada M, Cuchacovich RS, Shuler CL, Lin C-Y, BraunDK, Lee CH, Gladman DG. Ixekizumab, an interleukin-17A specific monoclonal antibody, for the treatment of biologic-naive patients with active psoriatic arthritis: results from the 24-week randomised, double-blind, placebo-controlled and active (adalimumab)-

controlled period of the phase III trial SPIRIT-P1.

Ann Rheum Dis 2017;76:79-86

2016

Jacquemin C, Rubio Vargas R, van den Berg R, Thévenin F, Lenczner G, Reijnierse M, Ferkal S, Le Corvoisier P, Rahmouni A, Loeuille D, Feydy A, Dougados M, [van der Heijde D](#), Claudepierre P. What is the reliability of non-trained investigators in recognising structural MRI lesions of sacroiliac joints in patients with recent inflammatory back pain? Results of the DESIR cohort.

RMD Open 2016;2(2):e000303

Haugen IK, Mathiessen A, Slatkowsky-Christensen B, Magnusson K, Bøyesen P, Sesseng S, [van der Heijde D](#), Kvien TK, Hammer HB. Synovitis and radiographic progression in non-erosive and erosive hand osteoarthritis: is erosive hand osteoarthritis a separate inflammatory phenotype?

Osteoarthritis Cartilage 2016;24:647-54

[van der Heijde D](#), Landewé RB, Mease PJ, McInnes IB, Conaghan PG, Pricop L, Ligozio G, Richards HB, Mpofu S. Secukinumab Provides Significant and Sustained Inhibition of Joint Structural Damage in a Phase III Study of Active Psoriatic Arthritis.

Arthritis Rheumatol 2016;68:1914-2

Aga A-B, Hammer HB, Olsen IC, Uhlig T, Kvien TK, [van der Heijde D](#), Fremstad H, Madland TM, Lexberg ÅS, Haukeland H, Rødevand E, Høili C, Stray H, Bendvold AN, Soldal DM, Bakland G, Lie E, Haavardsholm EA. First step in the development of an ultrasound joint inflammation score for rheumatoid arthritis using a data-driven approach.

Ann Rheum Dis 2016;75:1444-51

Dougados M, Demattei C, van den Berg R, Vo Hoang V, Thevenin F, Reijnierse M, Loeuille D, Feydy A, Claudepierre P, [van der Heijde D](#). Rate and Predisposing Factors for Sacroiliac Joint Radiographic Progression After a Two-Year Follow-up Period in Recent-Onset Spondyloarthritis.

Arthritis Rheumatol 2016;68:1904-13

Aga A-B, Berner Hammer H, Olsen IC, Uhlig T, Kvien TK, [van der Heijde D](#), Fremstad H, Madland TM, Lexberg ÅS, Haukeland H, Rødevand E, Høili C, Stray H, Noraas AL, Widding Hansen IJ, Bakland G, Lillegraven S, Lie E, Haavardsholm EA. Development of a feasible and responsive ultrasound inflammation score for rheumatoid arthritis through a data-driven approach

RMD Open 2016; 2:e000325, doi:10.1136/rmdopen-2016-000325

HuoY, Veldhuizen RD, [van der Heijde D](#), Besselink NJ, Jacobs JWG, van Laar JM, Viergever MA, Vincken KL, Lafeber FP, de Hair MJH, on behalf of the Society for Rheumatology Research Utrecht investigators. Automated joint space width quantification of hand and wrist joints: a proof of concept study.

Clin Exp Rheumatol 2016;34: S34-S39

Lambert RGW, Bakker PAC, [van der Heijde D](#), Weber U, Rudwaleit M, Hermann K-GA, Sieper J, Baraliakos X, Bennett A, Braun J, Burgos-Vargas R, Dougados M, Pedersen SJ, Jurik AG, Maksymowych WP, Marzo-Ortega H, Østergaard M,

Poddubnyy D, Reijnierse M, van den Bosch F, van der Horst-Bruinsma I, Landewé R. Defining active sacroiliitis on MRI for classification of axial spondyloarthritis: update by the ASAS MRI working group.

Ann Rheum Dis 2016;75:1958-63

Landewé RBM, Connell CA, Bradley JD, Wilkinson B, Gruben D, Strengholt S, van der Heijde D. Is radiographic progression in modern rheumatoid arthritis trials still a robust outcome? Experience from tofacitinib clinical trials.

Arthritis Research & Therapy 2016;18:212 DOI: 10.1186/s13075-016-1106-y

Huo Y, Vincken KL, van der Heijde D, De Hair MJH, Lafeber FP, Viergever MA. Automatic Quantification of Radiographic Finger Joint Space Width of Patients With Early Rheumatoid Arthritis.

IEEE Transactions on Biomedical Engineering 2016;63; 10.1109/TBME.2015.2512941

Machado PM, Baraliakos X, van der Heijde D, Braun J, Landewé R. MRI vertebral corner inflammation followed by fat deposition is the strongest contributor to the development of new bone at the same vertebral corner: a multilevel longitudinal analysis in patients with ankylosing spondylitis.

Ann Rheum Dis 2016;75:1486-93

Markusse IM, Landewé R, Wolterbeek R, Ho M, Jenkins M, van der Heijde D. Linear extrapolation of missing radiographic change scores in clinical trials does not spuriously overestimate group radiographic changes in rheumatoid arthritis.

Rheumatology 2016;55:1295-3000

Maksymowych WP, Dougados M, van der Heijde D, Sieper J, Braun J, Citera G, Van den Bosch F, Logeart I, Wajdula J, Jones H, Marshall L, Bonin R, Pedersen R, Vlahos B, Kotak S, Bukowski JF. Clinical and MRI responses to etanercept in early non-radiographic axial spondyloarthritis: 48-week results from the EMBARK study.

Ann Rheum Dis 2016;75:1328-35

de Hooge M, van den Berg R, Navarro-Compán V, Reijnierse M, van Gaalen F, Fagerli K, Landewé R, van Oosterhout M, Ramonda R, Huizinga T, van der Heijde D. Patients with chronic back pain of short duration from the SPACE cohort: which MRI structural lesions in the sacroiliac joints and inflammatory and structural lesions in the spine are most specific for axial spondyloarthritis?

Ann Rheum Dis 2016;75:1308-14

Conaghan PG, Østergaard M, Bowes MA, Wu C, Fuerst T, van der Heijde D, Irazoque-Palazuelos F, Soto-Raices O, Hrycaj P, Xie Z, Zhang R, Wyman BT, Bradley JD, Soma K, Wilkinson B. Comparing the effects of tofacitinib, methotrexate and the combination, on bone marrow oedema, synovitis and bone erosion in methotrexate-naive, early active rheumatoid arthritis: results of an exploratory randomised MRI study incorporating semiquantitative and quantitative techniques.

Ann Rheum Dis 2016;75:1024-33

Takeuchi T, Tanaka Y, Ishiguro N, Yamanaka H, Yoneda T, Ohira T, Okubo N, Genant HK, van der Heijde D. Effect of denosumab on Japanese patients with rheumatoid arthritis: a dose-response study of AMG 162 (Denosumab) in patients with Rheumatoid arthritis on methotrexate to Validate inhibitory effect on bone Erosion (DRIVE)—a 12-month, multicentre, randomised, double-blind, placebo-controlled, phase II clinical trial.

Ann Rheum Dis 2016;75:983-90

Nieuwenhuis WP, van Steenberg HW, Stomp W, Stijnen T, Huizinga TWJ, Bloem, [van der Heijde D](#), Reijnen M, van der Helm-van Mil AHM. The Course of Bone Marrow Edema in Early Undifferentiated Arthritis and Rheumatoid Arthritis: A Longitudinal Magnetic Resonance Imaging Study at Bone Level. *Arthritis Rheum* 2016;68:1080–8

[van der Heijde D](#), Landewé R. Are conventional radiographs still of value?
Curr Opin Rheum 2016;28:310-5

Navarro-Compán V, Ramiro S, Landewé R, Dougados M, Miceli-Richard C, Richette P, [van der Heijde D](#). Disease activity is longitudinally related to sacroiliac inflammation on MRI in male patients with axial spondyloarthritis: 2-years of the DESIR cohort.
Ann Rheum Dis 2016;75:874-8

de Bruin F, Ter Horst S, Bloem HL, van den Berg R, de Hooge M, van Gaalen F, Fagerli KM, Landewé R, van Oosterhout M, [van der Heijde D](#), Reijnen M. Prevalence of degenerative changes of the spine on magnetic resonance images and radiographs in patients aged 16-45 years with chronic back pain of short duration in the Spondyloarthritis Caught Early (SPACE) cohort.
Rheumatology 2016;55:56-65

de Bruin F, Ter Horst S, van den Berg R, de Hooge M, van Gaalen F, Fagerli KM, Landewé R, van Oosterhout M, Bloem JL, [van der Heijde D](#), Reijnen M. Signal intensity loss of the intervertebral discs in the cervical spine of young patients on fluid sensitive sequences.
Skeletal Radiol 2016;45:375-81

Gherghe AM, Ramiro S, Landewé R, Mihai C, [van der Heijde D](#). Association of the different types of radiographic damage with physical function in patients with rheumatoid arthritis: analysis of the RAPID trials.
RMD Open 2016;2:e000219.doi:10.1136/rmdopen-2015-000219

Kavanaugh A, [van der Heijde D](#), Beutler A, Gladman D, Mease P, Krueger GG, McInnes IB, Helliwell P, Coates LC, Xu S. Radiographic Progression of Patients With Psoriatic Arthritis Who Achieve Minimal Disease Activity in Response to Golimumab Therapy: Results Through 5 Years of a Randomized, Placebo-Controlled Study.
Arthritis Care Res 2016;68:267–74

Haugen IK, Slatkowsky Christensen B, Bøyesen P, Sesseng S, [van der Heijde D](#), Kvien TK. Increasing synovitis and bone marrow lesions are associated with incident joint tenderness in hand osteoarthritis.
Ann Rheum Dis 2016;75:702-8

Atsumi T, Yamamoto K, Takeuchi T, Yamanaka H, Ishiguro N, Tanaka Y, Eguchi K, Watanabe A, Origasa H, Yasuda S, Yamanishi Y, Kita Y, Matsubara T, Iwamoto M, Shoji T, Okada T, [van der Heijde D](#), Miyasaka N, Koike T. The first double-blind, randomised, parallel-group certolizumab pegol study in methotrexate-naive early rheumatoid arthritis patients with poor prognostic factors, C-OPERA, shows inhibition of radiographic progression.
Ann Rheum Dis 2016;75:75-83

Haugen IK, Slatkowsky-Christensen B, Bøyesen P, Sesseng S, [van der Heijde D](#), Kvien TK. MRI findings predict radiographic progression and development of erosions in hand osteoarthritis.

Ann Rheum Dis 2016;75:117-23

Inman RD, Baraliakos X, Hermann KA, Braun J, Deodhar A, van der Heijde D, Xu S, Hsu B. Serum biomarkers and changes in clinical/MRI evidence of golimumab-treated patients with ankylosing spondylitis: results of the randomized, placebo-controlled GO-RAISE study.

Arthritis Res Ther. 2016 Dec 28;18(1):304

2015

van den Berg R, de Hooge M, Bakker PA, van Gaalen F, Navarro-Compán V, Fagerli KM, Landewé R, van Oosterhout M, Ramonda R, Reijniere M, van der Heijde D. Metric Properties of the SPARCC Score of the Sacroiliac Joints - Data from Baseline, 3-month, and 12-month Followup in the SPACE Cohort.

J Rheumatol. 2015;42:1186-93

Kortekaas MC, Kwok WY, Reijniere M, Wolterbeek R, Bøyesen P, van der Heijde D, Bloem JL, Kloppenburg M. Magnetic Resonance Imaging in Hand Osteoarthritis: Intraobserver Reliability and Criterion Validity for Clinical and Structural Characteristics.

J Rheumatol 2015;42:1224-30

Takeuchi T, Tanaka Y, Ishiguro N, Yamanaka H, Yoneda T, Ohira T, Okubo N, Genant HK, van der Heijde D. Effect of denosumab on Japanese patients with rheumatoid arthritis: a dose–response study of AMG 162 (Denosumab) in patients with Rheumatoid arthritis on methotrexate to Validate inhibitory effect on bone Erosion (DRIVE)—a 12-month, multicentre, randomised, double-blind, placebo-controlled, phase II clinical trial.

Ann Rheum Dis Published 19 November 2015, doi:10.1136/annrheumdis-2015-208052

Huo Y, De Hair MJH, Shaib YO, van der Heijde D, Kuchuk NO, Viergever MA, van Laar JM, Vincken KL, FP Lafeber. Computerised versus conventional methodology of radiographic joint destruction assessment in early rheumatoid arthritis.

RMD Open 2015; 1:e000148, doi:10.1136/rmdopen-2015-000148

Haugen IK, Slatkowsky Christensen B, Bøyesen P, Sesseng S, van der Heijde D, Kvien TK. Increasing synovitis and bone marrow lesions are associated with incident joint tenderness in hand osteoarthritis.

Ann Rheum Dis 2015;75:117-23

Atsumi T, Yamamoto K, Takeuchi T, Yamanaka H, Ishiguro N, Tanaka Y, Eguchi K, Watanabe A, Origasa H, Yasuda S, Yamanishi Y, Kita Y, Matsubara T, Iwamoto M, Shoji T, Okada T, van der Heijde D, Miyasaka N, Koike T. The first double-blind, randomised, parallel-group certolizumab pegol study in methotrexate-naive early rheumatoid arthritis patients with poor prognostic factors, C-OPERA, shows inhibition of radiographic progression. Danila MI, Westfall AO, Raman K,

Chen L, Reynolds RJ, Hughes LB, Arnett DK, McGwin G, Szalai AJ, van der Heijde DM, Conn D, Callahan LF, Moreland LW, Bridges Jr SL. The role of genetic variants in CRP in radiographic severity in African Americans with early and established rheumatoid arthritis.

Genes and Immunity 2015;16:446-51

Sieper J, van der Heijde D, Dougados M, Maksymowych WP, Scott BB, Boice JA, Berd Y, Bergman G, Curtis S, Tzontcheva A,

Huyck S, Weng HH. A Randomized, Double-Blind, Placebo-Controlled, Sixteen-Week Study of Subcutaneous Golimumab in Patients With Active Nonradiographic Axial Spondyloarthritis.

Arthritis Rheum 2015;67:2702–12

van den Berg R, Lenczner G, Thévenin F, Claudepierre P, Feydy A, Reijnierse M, Saraux A, Rahmouni A, Dougados M, van der Heijde D. Classification of axial SpA based on positive imaging (radiographs and/or MRI of the sacroiliac joints) by local rheumatologists or radiologists versus central trained readers in the DESIR cohort.

Ann Rheum Dis 2015;74:2016-21

Navarro-Compán V, de Miguel E, van der Heijde D, Landewé R, Almodóvar R, Montilla C, Beltrán E, Zarco P. Spondyloarthritis features forecasting the presence of HLA-B27 or sacroiliitis on magnetic resonance imaging in patients with suspected axial spondyloarthritis: Results from a cross-sectional study in the ESPeranza Cohort.

Arthritis Res Ther 2015;17:265

Mease PJ, McInnes IB, Kirkham B, Kavanaugh A, Rahman P, van der Heijde D, Landewé R, Nash P, Pricop L, Yuan J, Richards HB, Mpofo S, for the FUTURE 1 Study Group. Secukinumab Inhibition of Interleukin-17A in Patients with Psoriatic Arthritis. N Engl J Med 2015;373:1329-39

Smolen JS, van Vollenhoven R, Kavanaugh A, Strand V, Vencovsky J, Schiff M, Landewé R, Haraoui B, Arendt C, Mountian I, Carter D, van der Heijde D. Certolizumab pegol plus methotrexate 5-year results from the rheumatoid arthritis prevention of structural damage (RAPID) 2 randomized controlled trial and long-term extension in rheumatoid arthritis patients.

Arthritis Res Ther 2015;17:art no 245

Ramiro S, Landewé R, van Tubergen A, Boonen A, Stolwijk C, Dougados M, van den Bosch F, van der Heijde D. Lifestyle factors may modify the effect of disease activity on radiographic progression in patients with ankylosing spondylitis: a longitudinal analysis.

RMD Open 2015;1:e000153, doi:10.1136/rmdopen-2015-000153

Mangnus L, van Steenbergen HW, Lindqvist E, Brouwer E, Reijnierse M, Huizinga TWJ, Gregersen PK, Berglin E, Rantapää-Dahlqvist S, van der Heijde D, van der Helm-van Mil, A.H.M. Studies on ageing and the severity of radiographic joint damage in rheumatoid arthritis.

Arthritis Res Ther 2015;17:art no 222

Barnabe C, Sun Y, Boire G, Hitchon CA, Haraoui B, Carter Thorne J, Tin D, van der Heijde D, Curtis JR, Jamal S, Pope JE, Keystone EC, Bartlett S, Bykerk VP, CATCH Investigators. Heterogeneous Disease Trajectories Explain Variable Radiographic, Function and Quality of Life Outcomes in the Canadian Early Arthritis Cohort (CATCH).

PLoS ONE 10(8): e0135327

Magnusson K, Slatkowsky-Christensen B, van der Heijde D, Kvien TK, Hagen KB, Haugen IK. Body mass index and progressive hand osteoarthritis: Data from the Oslo hand osteoarthritis cohort.

Scand J Rheumatol 2015;44:331-6

Mease P, Deodhar A, Fleischmann R, Wollenhaupt J, Gladman D, Leszczyński P, Vitek P, Turkiewicz A, Khraishi M, FitzGerald O, Landewé R, de Longueville M, Hoepken B, Peterson L, van der Heijde D. Effect of certolizumab pegol over 96 weeks in

patients with psoriatic arthritis with and without prior antitumour necrosis factor exposure.
RMD Open 2015;1:e000119.doi:10.1136/rmdopen-2015-000119

Mandl P, Navarro-Compán V, Terslev L, Aegerter P, [van der Heijde D](#), D'Agostino MA, Baraliakos X, Pedersen SJ, Jurik AG, Naredo E, Schueller-Weidekamm C, Weber U, Wick MC, Bakker PAC, Filippucci E, Conaghan PG, Rudwaleit M, Schett G, Sieper J, Tarp S, Marzo-Ortega H, Østergaard M. EULAR recommendations for the use of imaging in the diagnosis and management of spondyloarthritis in clinical practice.
Ann Rheum Dis 2015;74:1327-39

Genovese MC, Fleischmann R, Kivitz AJ, Rell-Bakalarska M, Martincova R, Fiore S, Rohane P, van Hoogstraten H, Garg A, Fan C, van Adelsberg J, Weinstein SP, Graham NMH, Stahl N, Yancopoulos G, Huizinga TWJ, [van der Heijde D](#). Sarilumab Plus Methotrexate in Patients With Active Rheumatoid Arthritis and Inadequate Response to Methotrexate: Results of a Phase III Study.
Arthritis Rheumatol 2015;67:1424–37

Landewé R, Smolen JS, Florentinus S, Chen S, Guérette B, [van der Heijde D](#). Existing joint erosions increase the risk of joint space narrowing independently of clinical synovitis in patients with early rheumatoid arthritis.
Arthritis Res Ther 2015;17:133

Navarro-Compán V, Gherghe AM, Smolen JS, Aletaha D, Landewé R, [van der Heijde D](#). Relationship between disease activity indices and their individual components and radiographic progression in RA: a systematic literature review.
Rheumatology 2015;54:83-90

Hemke R, Nusman CM, [van der Heijde DMFM](#), Doria AS, Kuijpers TW, Maas M, van Rossum MAJ. Frequency of joint involvement in juvenile idiopathic arthritis during a 5-year follow-up of newly diagnosed patients: implications for MR imaging as outcome measure.
Rheumatol Int 2015;35:351-7

Stomp W, Krabben A, [van der Heijde D](#), Huizinga TWJ, Bloem JL, Østergaard M, van der Helm-van Mil, AHM, Reijnen M. Aiming for a simpler early arthritis MRI protocol: can Gd contrast administration be eliminated?
European Radiology 2015;25:1520-7

Nieuwenhuis WP, Krabben A, Stomp W, Huizinga TWJ, [van der Heijde D](#), Bloem JL, van der Helm-van Mil AHM, Reijnen M. Evaluation of Magnetic Resonance Imaging–Detected Tenosynovitis in the Hand and Wrist in Early Arthritis.
Arthritis Rheumatol 2015;67:869–76

van Onna M, van Tubergen A, Jurik A, [van der Heijde D](#), Landewé R. Natural course of bone marrow oedema on magnetic resonance imaging of the sacroiliac joints in patients with early inflammatory back pain: a 2-year follow-up study.
Scand J Rheumatol 2015;44:129-34

Sieper J, Landewé R, Rudwaleit M, [van der Heijde D](#), Dougados M, Mease PJ, Braun J, Deodhar A, Kivitz A, Walsh J, Hoepken B, Nurminen T, Maksymowych WP. Effect of Certolizumab Pegol over 96 Weeks in Patients with Axial Spondyloarthritis: Results from a Phase 3 Randomized Trial.
Arthritis Rheumatol 2015;67:668-77

Deodhar A, Braun J, Inman RD, van der Heijde D, Zhou Y, Xu S, Han C, Hsu B. Golimumab administered subcutaneously every 4 weeks in ankylosing spondylitis: 5-year results of the GO-RAISE study.

Ann Rheum Dis 2015;74:757-61

Moltó A, Paternotte S, van der Heijde D, Claudepierre P, Rudwaleit M, Dougados M. Evaluation of the validity of the different arms of the ASAS set of criteria for axial spondyloarthritis and description of the different imaging abnormalities suggestive of spondyloarthritis: data from the DESIR cohort.

Ann Rheum Dis 2015;74:746-51

Krabben A, Stomp W, Huizinga TWJ, van der Heijde D, Bloem JL, Reijniere M, van der Helm-van Mil AHM. Concordance between inflammation at physical examination and on MRI in patients with early arthritis.

Ann Rheum Dis 2015; 74:506-512

Landewé R, Østergaard M, Keystone EC, Florentinus S, Liu S, van der Heijde D. Analysis of integrated radiographic data from two long-term, open-label extension studies of adalimumab for the treatment of rheumatoid arthritis.

Arthritis Care Res 2015;2:108-6

Navarro-Compán V, Landewé R, Provan SA, Ødegard S, Uhlig T, Kvien TK, Keszei AP, Ramiro S, van der Heijde D. Relationship between types of radiographic damage and disability in patients with rheumatoid arthritis in the EURIDISS cohort: a longitudinal study.

Rheumatology 2015;54:83-90

Ramiro S, Stolwijk C, van Tubergen A, van der Heijde D, Dougados M, van den Bosch F, Landewé R. Evolution of radiographic damage in ankylosing spondylitis: a 12 year prospective follow-up of the OASIS study.

Ann Rheum Dis 2015;74:52-9

2014

Haugen IK, Slatkowsky-Christensen B, Bøyesen P, Sesseng S, van der Heijde D, Kvien TK. MRI findings predict radiographic progression and development of erosions in hand osteoarthritis.

Ann Rheum Dis Published 9 September 2014, doi:10.1136/annrheumdis-2014-205949

Stomp W, Krabben A, van der Heijde D, Huizinga TW, Bloem JL, van der Helm-van Mil AH, Reijniere M. Are rheumatoid arthritis patients discernible from other early arthritis patients using 1.5T extremity magnetic resonance imaging? A large cross-sectional study.

J Rheumatol 2014;41:1630-7

Ramiro S, van der Heijde D, van Tubergen A, Stolwijk C, Dougados M, van den Bosch F, Landewé R. Higher disease activity leads to more structural damage in the spine in ankylosing spondylitis: 12-year longitudinal data from the OASIS cohort.

Ann Rheum Dis. 2014;73:1455-61

Hemke R, Doria AS, Tzaribachev N, Maas M, van der Heijde DM, van Rossum MA. Selecting Magnetic Resonance Imaging (MRI) Outcome Measures for Juvenile Idiopathic Arthritis (JIA) Clinical Trials: First Report of the MRI in JIA Special Interest Group.

J Rheumatol 2014;41:354-8

Baraliakos X, Heldmann F, Callhoff J, Listing J, Appelboom T, Brandt J, Van den Bosch F, Breban M, Burmester GR, Dougados M, Emery P, Gaston H, Grunke M, van der Horst-Bruinsma IE, Landewé R, Leirisalo-Repo M, Sieper J, De Vlam K, Pappas D, Kiltz U, van der Heijde D, Braun J. Which spinal lesions are associated with new bone formation in patients with ankylosing spondylitis treated with anti-TNF agents? A long-term observational study using MRI and conventional radiography.

Ann Rheum Dis 2014;73:1819-25

Nieuwenhuis WP, van Nies JA, Reijnierse M, van der Heijde D, van der Helm-Van Mil AH. Is the modified disease activity score superior to the disease activity score in early arthritis and rheumatoid arthritis? Comment on the article by Baker et al.

Arthritis Rheumatol. 2014;66:3244-5

Visser AW, Bøyesen P, Haugen IK, Schoones JW, van der Heijde DM, Rosendaal FR, Kloppenburg M. Radiographic scoring methods in hand osteoarthritis - a systematic literature search and descriptive review.

Osteoarthritis Cartilage 2014;22:1710-23

Keystone E, Landewé, van Vollenhoven R, Combe B, Strand V, Mease P, Shaughnessy L, VanLunen B, van der Heijde D. Long-term safety and efficacy of certolizumab pegol in combination with methotrexate in the treatment of rheumatoid arthritis: 5-year results from the RAPID 1 trial and open-label extension.

Ann Rheum Dis 2014; 73:2094-100

Krabben A, Stomp W, van Nies JAB, Huizinga TWJ, van der Heijde D, Bloem JL, Reijnierse M, van der Helm-van Mil AHM. MRI-detected subclinical joint inflammation is associated with radiographic progression.

Ann Rheum Dis 2014;73:2034-7

Ramiro S, van Tubergen A, van der Heijde D, Stolwijk C, Bookelman G, Dougados M, van den Bosch F, Landewé RBM. Erosions and sclerosis on radiographs precede the subsequent development of syndesmophytes at the same site: a twelve-year prospective followup of patients with ankylosing spondylitis.

Arthritis Rheumatology 2014;66:2773–2779

van den Berg R, Lenczner G, Feydy A, van der Heijde D, Reijnierse M, Saraux A, Rahmouni A, Dougados M, Claudepierre P. Reading of sacroiliac joints on plain pelvic radiographs: Agreement between clinical practice and trained central reading. Results of the DESIR-cohort.

Arthritis Rheumatol 2014;66:2403-11

Stomp W, Krabben A, van der Heijde D, Huizinga TWJ, Bloem JL, van der Helm-van Mil AHM, Reijnierse M. Aiming for a shorter rheumatoid arthritis MRI protocol: can contrast-enhanced MRI replace T2 for the detection of bone marrow oedema?

Eur Radiol 2014;24:2614-22

Yamamoto K, Takeuchi T, Yamanaka H, Ishiguro N, Tanaka Y, Eguchi K, Watanabe A, Origasa H, Iwai K, Sakamaki Y, van der Heijde D, Miyasaka N, Koike T. Efficacy and safety of certolizumab pegol without methotrexate co-administration in

Japanese patients with active rheumatoid arthritis: the HIKARI randomized, placebo-controlled trial.
Mod Rheumatol 2014;24:715-24

Hafström I, Engvall I-L, Rönnelid J, Boonen A, [van der Heijde D](#), Svensson B. Rheumatoid factor and anti-CCP do not predict progressive joint damage in patients with early rheumatoid arthritis treated with prednisolone: A randomised study.
BMJ Open 2014;4:e005246

Kavanaugh A, McInnes IB, Mease P, Krueger GG, Gladman D, [van der Heijde D](#), Zhou Y, Lu J, Leu JH, Goldstein N, Beutler A. Clinical efficacy, radiographic and safety findings through 5 years of subcutaneous golimumab treatment in patients with active psoriatic arthritis: results from a long-term extension of a randomised, placebo-controlled trial, the GO-REVEAL study).
Ann Rheum Dis 2014;73:1689-94

Dougados M, [van der Heijde D](#), Sieper J, Braun J, Maksymowych WP, Citera G, Miceli-Richard C, Wei JC, Pedersen R, Bonin R, Rahman MU, Logeart I, Wajdula J, Koenig AS, Vlahos B, Alvarez D, Bukowski JF. Symptomatic efficacy of etanercept and its effects on objective signs of inflammation in early nonradiographic axial spondyloarthritis: a multicenter, randomized, double-blind, placebo-controlled trial.
Arthritis Rheumatol. 2014 Aug;66(8):2091-102

van Gaalen FA, Bakker PAC, de Hooge M, Schoones JW, [van der Heijde D](#). Assessment of sacroiliitis by radiographs and MRI: Where are we now?
Curr Opin Rheum 2014;26:384-8

van Onna M, van Tubergen A, [van der Heijde DM](#), Jurik AG, Landewé R. Bone Marrow Edema on Magnetic Resonance Imaging (MRI) of the Sacroiliac Joints Is Associated with Development of Fatty Lesions on MRI over a 1-year Interval in Patients with Early Inflammatory Low Back Pain: A 2-year Followup Study.
J Rheumatol 2014;41:1088-94

de Rooy DPC, van Nies JAB, Kapetanovic MC, Kristjansdottir H, Andersson MLE, Forslind K, [van der Heijde DMFM](#), Gregersen PK, Lindqvist E, Huizinga TWJ, Gröndal G, Svensson B, van der Helm-van Mil AHM. Smoking as a risk factor for the radiological severity of rheumatoid arthritis: a study on six cohorts.
Ann Rheum Dis 2014; 73:1384-7

Maksymowych WP, [van der Heijde D](#), Allaart CF, Landewé R, Boire G, Tak PP, Gui Y, Ghahary A, Kilani R, Marotta A. 14-3-3 η is a novel mediator associated with the pathogenesis of rheumatoid arthritis and joint damage.
Arthritis Res Ther 2014;16:R99

Braun J, Baraliakos X, Hermann K-GA, Deodhar A, [van der Heijde D](#), Inman R, Beutler A, Zhou Y, Xu S, Hsu B. The effect of two golimumab doses on radiographic progression in ankylosing spondylitis: results through 4 years of the GO-RAISE trial.
Ann Rheum Dis 2014;73:1107-13

Navarro-Compán V, [van der Heijde D](#), Ahmad HA, Miller CG, Wolterbeek R, Landewé R. Measurement error in the assessment of radiographic progression in rheumatoid arthritis (RA) clinical trials: the smallest detectable change (SDC) revisited.

Ann Rheum Dis 2014;73:1067-70

Nam JL, Villeneuve E, Hensor EMA, Wakefield RJ, Conaghan PG, Green MJ, Gough A, Quinn M, Reece R, Cox SR, Buch MH, van der Heijde DM, Emery P. A randomised controlled trial of etanercept and methotrexate to induce remission in early inflammatory arthritis: the EMPIRE trial.

Ann Rheum Dis 2014;73:1027-36

van Onna M, van Tubergen A, van der Heijde D, Jurik AG, Landewé R. Gadolinium contrast-enhanced MRI sequence does not have an incremental value in the assessment of sacroiliitis in patients with early inflammatory back pain by using MRI in combination with pelvic radiographs: a 2-year follow-up study.

Clin Exp Rheumatol 2014;32:225-30

van der Heijde D, Sieper J, Maksymowych W, Brown M, Lambert R, Rathmann S, Pangal A. Spinal inflammation in the absence of sacroiliac joint inflammation on MRI in patients with active non-radiographic axial spondyloarthritis.

Arthritis Rheum 2014;66:667-73

Hemke R, Doria AS, Tzaribachev N, Maas M, van der Heijde DM, van Rossum MA. Selecting Magnetic Resonance Imaging (MRI) Outcome Measures for Juvenile Idiopathic Arthritis (JIA) Clinical Trials: First Report of the MRI in JIA Special Interest Group.

J Rheumatol 2014;41:354-8

Kroon FPB, van der Burg LRA, Ramiro S, Landewé RBM, Buchbinder R, van der Heijde D. Non-steroidal anti-inflammatory drugs (NSAIDs) for axial spondyloarthritis (ankylosing spondylitis and non-radiographic axial spondyloarthritis) (Protocol). Cochrane Database of Systematic Reviews 2014;1:CD010952. DOI: 10.1002/14651858.CD010952

van Tuyl LHD, van der Heijde D, Knol DL, Boers M. Chronological reading of radiographs in rheumatoid arthritis increases efficiency and does not lead to bias.

Ann Rheum Dis 2014;73:391-5

Landewé R, Braun J, Deodhar A, Dougados M, Maksymowych WP, Mease PJ, Reveille JD, Rudwaleit M, van der Heijde D, Stach C, Hoepken B, Fichtner A, Coteur G, de Longueville M, Sieper J. Efficacy of Certolizumab Pegol on Signs and Symptoms of Axial Spondyloarthritis including Ankylosing Spondylitis: 24 Week Results of a Double-Blind Randomized Placebo-Controlled Phase 3 Study.

Ann Rheum Dis 2014;73:39-47

Schiff M, Weinblatt ME, Valente R, van der Heijde D, Citera G, Elegbe A, Maldonado M, Fleischmann R. Head-to-head comparison of subcutaneous abatacept versus adalimumab for rheumatoid arthritis: two-year efficacy and safety findings from AMPLE trial.

Ann Rheum Dis 2014;73:86-94

Mease PJ, Fleischmann R, Deodhar AA, Wollenhaupt J, Khraishi M, Kielar D, Woltering F, Stach C, Hoepken B, Arledge T, van der Heijde D. Effect of certolizumab pegol on signs and symptoms in patients with psoriatic arthritis: 24-week results of a Phase 3 double-blind randomised placebo-controlled study (RAPID-PsA).

Ann Rheum Dis 2014;73:48-55

van der Heijde D, Fleischmann R, Wollenhaupt J, Deodhar AA, Kielar D, Woltering F, Stach C, Hoepken B, Arledge T Mease PJ. Effect of different imputation approaches on the evaluation of radiographic progression in patients with psoriatic arthritis: results of the RAPID-PsA 24-week phase III double-blind randomised placebo-controlled study of certolizumab pegol.
Ann Rheum Dis 2014;73:233-7

Nam JL, Villeneuve E, Hensor EMA, Conaghan PG, Keen HI, Buch MH, Gough AK, Green MJ, Helliwell PS, Keenan AM, Morgan AW, Quinn M, Reece R, van der Heijde DM, Wakefield RJ, Emery P. Remission induction comparing infliximab and high-dose intravenous steroid, followed by treat-to-target: a double-blind, randomised, controlled trial in new-onset, treatment-naive, rheumatoid arthritis (the IDEA study).
Ann Rheum Dis 2014;73:75-85

Knevel R, Tsonaka R, Le Cessie S, van der Linden M, Huizinga T, van der Heijde D, Houwing-Duistermaat J, van der Helm-van Mil A. Comparison of methodologies for analysing the progression of joint destruction in rheumatoid arthritis.
Scand J Rheumatol 2014;42:182-9

Keystone EC, van der Heijde D, Kavanaugh A, Kupper H, Liu S, Guérette B, Mozaffarian N. Clinical, functional, and radiographic benefits fo longterm adalimumab plus methotrexate: final 10-year data in longstanding rheumatoid arthritis.
J Rheumatol 2014;40:1487-97

Haugen I, Slatkowsky-Christensen B, Bøyesen P, van der Heijde D, Kvien T. Cross-sectional and longitudinal associations between radiographic features and measures of pain and physical function in hand osteoarthritis.
Osteoarthritis Cartilage 2014;21:1191-8

2013

Yamamoto K, Takeuchi T, Yamanaka H, Ishiguro N, Tanaka Y, Eguchi K, Watanabe A, Origasa H, Iwai K, Sakamaki Y, van der Heijde D, Miyasaka N, Koike T. Efficacy and safety of certolizumab pegol without methotrexate co-adminstration in Japanese patients with active rheumatoid arthritis: the HIKARI randomized, placebo-controlled trial.
Mod Rheumatol 2013 published November 1

van der Heijde D, Sieper J, Maksymowych W, Brown M, Lambert R, Rathmann S, Pangal A. Spinal inflammation in the absence of sacroiliac joint inflammation on MRI in patients with active non-radiographic axial spondyloarthritis.
Arthritis Rheum 2013 published November 18

Keystone E, Breedveld F, van der Heijde D, Landewé R, Florentinus S, Arulmani U, Liu S, Kupper H, Kavanaugh A. Longterm effect of delaying combination therapy with tumor necrosis factor inhibitor in patients with aggressive early rheumatoid arthritis: 10-year efficacy and safety of adalimumab from the randomized controlled PREMIER trial with open-label extension.
J Rheumatol 2013 published November 15

Ramiro S, Stolwijk C, van Tubergen A, van der Heijde D, Dougados M, van den Bosch F, Landewé R. Evolution of radiographic damage in ankylosing spondylitis: a 12 year prospective follow-up of the OASIS study.
Ann Rheum Dis published 16 August 2013, 10.1136/annrheumdis-2013-204055

Keystone E, Landewé, van Vollenhoven R, Combe B, Strand V, Mease P, Shaughnessy L, VanLunen B, [van der Heijde D](#). Long-term safety and efficacy of certolizumab pegol in combination with methotrexate in the treatment of rheumatoid arthritis: 5-year results from the RAPID 1 trial and open-label extension. *Ann Rheum Dis* published 5 August 2013, 10.1136/annrheumdis-2013-203695

Baraliakos X, Heldmann F, Callhoff J, Listing J, Appelboom T, Brandt J, Van den Bosch F, Breban M, Burmester GR, Dougados M, Emery P, Gaston H, Grunke M, van der Horst-Bruinsma IE, Landewé R, Leirisalo-Repo M, Sieper J, De Vlam K, Pappas D, Kiltz U, [van der Heijde D](#), Braun J. Which spinal lesions are associated with new bone formation in patients with ankylosing spondylitis treated with anti-TNF agents? A long-term observational study using MRI and conventional radiography. *Ann Rheum Dis* published 14 July 2013, 10.1136/annrheumdis-2013-203425

Braun J, Baraliakos X, Hermann K-GA, Deodhar A, [van der Heijde D](#), Inman R, Beutler A, Zhou Y, Xu S, Hsu B. The effect of two golimumab doses on radiographic progression in ankylosing spondylitis: results through 4 years of the GO-RAISE trial. *Ann Rheum Dis* published 3 May 2013, 10.1136/annrheumdis-2012-203075

Navarro-Compan V, [van der Heijde D](#), Ahmad HA, Miller CG, Wolterbeek R, Landewé R. Measurement error in the assessment of radiographic progression in rheumatoid arthritis (RA) clinical trials: the smallest detectable change (SDC) revisited. *Ann Rheum Dis* published 20 April 2013, 10.1136/annrheumdis-2012-202939

van Tuyl LHD, [van der Heijde D](#), Knol DL, Boers M. Chronological reading of radiographs in rheumatoid arthritis increases efficiency and does not lead to bias. *Ann Rheum Dis* published 24 January 2013, 10.1136/annrheumdis-2012-202876

Hemke R, Doria AS, Tzaribachev N, Maas M, [van der Heijde DM](#), van Rossum MA. Selecting Magnetic Resonance Imaging (MRI) Outcome Measures for Juvenile Idiopathic Arthritis (JIA) Clinical Trials: First Report of the MRI in JIA Special Interest Group. *J Rheumatol* 2013 Nov 1;

Ciurea A, Scherer A, Exer P, Bernhard J, Dudler J, Beyeler B, Kissling R, Stekhoven D, Rufibach K, Tamborrini G, Weiss B, Mueller R, Nissen M, Michel B, [van der Heijde D](#), Dougados M, Boonen A, Weber U; Rheumatologists of the Swiss Clinical Quality Management Program for Axial Spondyloarthritis. Tumor necrosis factor-alpha inhibition in radiographic and non-radiographic axial spondyloarthritis: results from a large observational cohort. *Arthritis Rheum* 2013 Aug 27, doi: 10.1002/art38140

Grandaunet B, Watterdal Syversen S, Hoff M, Haugeberg G, [van der Heijde D](#), Kvien TK, Standal T. Dickkopf-1 Is Associated with Periarticular Bone Loss in Patients with Rheumatoid Arthritis. *OJRA* 2013;3:216-20

Kavanaugh A, McInnes IB, Mease PJ, Krueger GG, Gladman DD, [van der Heijde D](#), Mudivarthy S, Xu W, Mack M, Xu Z, Beutler A. Clinical efficacy, radiographic and safety findings through 2 years of golimumab treatment in patients with active psoriatic arthritis: results from a long-term extension of the randomised, placebo-controlled GO-REVEAL study.

Ann Rheum Dis 2013;72:1777-85

Siebuhr A, Bay-Jensen A, Leeming DJ, Platt A, Byrjalsen I, Christiansen C, van der Heijde D, Karsdal M. Serological identification of fast progressors of structural damage with rheumatoid arthritis. Arthritis Research & Therapy 2013, 15:R86

Krabben A, Stomp W, van der Heijde DMFM, van Nies JAB, Bloem JL, Huizinga TWJ, Reijnen M, van der Helm-van Mil AHM. MRI of hand and foot joints of patients with anticitrullinated peptide antibody positive arthralgia without clinical arthritis. Ann Rheum Dis 2013;72:1540-4

Navarro-Compán V, Landewé R, Ahmad HA, Miller CG, Xu D, Wolterbeek R, van der Heijde D. Rate of adjudication of radiological progression in rheumatoid arthritis randomized controlled trials depending on preset limits of agreement: a pooled analysis from 15 randomized trials. Rheumatology 2013;52:1404-7

de Hooge M, van den Berg R, Navarro-Compán V, van Gaalen F, van der Heijde D, Huizinga T, Reijnen M. Magnetic resonance imaging of the sacroiliac joints in the early detection of spondyloarthritis: No added value of gadolinium compared with short tau inversion recovery sequence. Rheumatology 2013;52:1220-4

Landewé R, Strand V, van der Heijde D. From inhibition of radiographic progression to maintaining structural integrity: a methodological framework for radiographic progression in rheumatoid arthritis and psoriatic arthritis clinical trials. Ann Rheum Dis 2013;72:1113-7

Mandl P, Balint PV, Brault Y, Backhaus M, D'Agostino MA, Grassi W, van der Heijde D, de Miguel E, Wakefield RJ, Logeart I, Dougados M. Clinical and Ultrasound-Based Composite Disease Activity Indices in Rheumatoid Arthritis: Results From a Multicenter, Randomized Study. Arthritis Care Res 2013;65:879-87

Smolen JS, van der Heijde DM, Keystone EC, van Vollenhoven RF, Goldring MB, Guerette B, Cifaldi MA, Chen N, Liu S, Landewé RBM. Association of joint space narrowing with impairment of physical function and work ability in patients with early rheumatoid arthritis: protection beyond disease control by adalimumab plus methotrexate. Ann Rheum Dis 2013;72:1156-62

Koevoets R, Dirven L, Klarenbeek NB, van Krugten MV, Roday HK, van der Heijde DMFM, Huizinga TWJ, Kerstens PSJM, Lems WF, Allaart CF. Insights in the relationship of joint space narrowing versus erosive joint damage and physical functioning of patients with RA. Ann Rheum Dis 2013;72:870-4

Sieper J, van der Heijde D, Dougados M, Mease PJ, Maksymowych WP, Brown MA, Arora V, Pangan AL. Efficacy and safety of adalimumab in patients with non-radiographic axial spondyloarthritis: results of a randomised placebo-controlled trial (ABILITY-1). Ann Rheum Dis 2013;72:815-22

Colebatch AN, Edwards CJ, Østergaard M, [van der Heijde D](#), Balint PV, D'Agostino M-A, Forslind K, Grassi W, Haavardsholm EA, Haugeberg G, Jurik A-G, Landewé RBM, Naredo E, O'Connor PJ, Ostendorf B, Potocki K, Schmidt WA, Smolen JS, Sokolovic S, Watt I, Conaghan PG. EULAR recommendations for the use of imaging of the joints in the clinical management of rheumatoid arthritis.
Ann Rheum Dis 2013;72:804-14

Vossen MHE, den Broeder AA, Hendriks-Roelofs F, [van der Heijde DMFM](#), Reijnen M. Improvement in deployment of MRI of the sacroiliac joints in patients suspected for spondyloarthritis using a targeted intervention: A case study.
Rheumatology 2013;52:933-8

Maehlen, M.T., Provan, S.A., de Rooy, D.P.C., van der Helm-van Mil, A.H.M., Krabben, A., Saxne, T., Lindqvist, E., Semb, A.G., Uhlig, T., van der Heijde, D., Mero, I.L., Olsen, I.C., Kvien, T.K., Lie, B.A. Associations between APOE Genotypes and Disease Susceptibility, Joint Damage and Lipid Levels in Patients with Rheumatoid Arthritis.
PLoS ONE , 2013;8:art. no. e60970

Engvall I-L, Svensson B, Boonen A, [van der Heijde D](#), Lerner UH, Hafström I. Low-dose prednisolone in early rheumatoid arthritis inhibits collagen type I degradation by matrix metalloproteinases as assessed by serum 1CTP-A possible mechanism for specific inhibition of radiological destruction.
Rheumatology 2013;52:733-42

Dougados M, Devauchelle-Pensec V, Ferlet JF, Jousse-Joulin S, D'Agostino M-A, Backhaus M, Bentin J, Chales G, Chary-Valckenaere I, Conaghan P, Wakefield RJ, Etcheparre F, Gaudin P, Grassi W, [van der Heijde D](#), Mariette X, Naredo E, Szkudlarek M. The ability of synovitis to predict structural damage in rheumatoid arthritis: a comparative study between clinical examination and ultrasound.
Ann Rheum Dis 2013;72:665-71

[van der Heijde D](#), Tanaka Y, Fleischmann R, Keystone E, Kremer J, Zerbini C, Cardiel MH, Cohen S, Nash P, Song Y-W, Tegzová D, Wyman BT, Gruben D, Benda B, Wallenstein G, Krishnaswami S, Zwillich SH, Bradley JD, Connell CA, and the ORAL Scan Investigators. Tofacitinib (CP-690,550) in patients with rheumatoid arthritis receiving methotrexate: Twelve-month data from a twenty-four-month phase III randomized radiographic study.
Arthritis Rheum 2013;65:559-70

Ramiro S, van Tubergen A, Stolwijk C, Landewe R, van den Bosch F, Dougados M, [van der Heijde D](#). Scoring radiographic progression in ankylosing spondylitis: Should we use the modified Stoke Ankylosing Spondylitis Spine Score (mSASSS) or the Radiographic Ankylosing Spondylitis Spinal Score (RASSS)?
Arthritis Research & Therapy 2013, 15:R14 (17 January 2013)

Weinblatt ME, Schiff M, Valente R, [van der Heijde D](#), Citera G, Zhao C, Maldonado M, Fleischmann R. Head-to-Head Comparison of Subcutaneous Abatacept versus Adalimumab for Rheumatoid Arthritis.
Arthritis Rheum 2013;65:28-38

2012

Machado P, Landewé RBM, Braun J, Baraliakos X, Hermann K-GA, Hsu B, Baker D, [van der Heijde D](#). MRI inflammation and its relation with measures of clinical disease activity and different treatment responses in patients with ankylosing spondylitis treated with a tumour necrosis factor inhibitor.

Ann Rheum Dis 2012;71:2002-5

van Tubergen A, [van der Heijde D](#), Dougados M, Mielants H, Landewé R. Are syndesmophytes most prevalent in the lumbar or in the cervical spine in patients with ankylosing spondylitis and do they develop in a specific direction?

Rheumatology 2012;51:1432-9

Keystone EC, Combe B, Smolen J, Strand V, Goel N, van Vollenhoven R, Mease P, Landewé R, Fleischmann R, Luijckens K, [van der Heijde D](#). Sustained efficacy of certolizumab pegol added to methotrexate in the treatment of rheumatoid arthritis: 2-year results from the RAPID 1 trial.

Rheumatology 2012;51:1628-38

Kroon F, Landewé R, Dougados M, [van der Heijde D](#). Continuous NSAID use reverts the effects of inflammation on radiographic progression in patients with ankylosing spondylitis.

Ann Rheum Dis 2012;71:1623-9

Moreland LW, O'Dell JR, Paulus HE, Curtis JR, Bathon JM, St.Clair EW, Bridges Jr SL, Zhang J, McVie T, Howard G, [van der Heijde D](#), Cofield SS. A randomized comparative effectiveness study of oral triple therapy versus etanercept plus methotrexate in early aggressive rheumatoid arthritis: The treatment of early aggressive rheumatoid arthritis trial.

Arthritis Rheum 2012;64:2824-35

Kavanaugh A, [van der Heijde D](#), McInnes IB, Mease P, Krueger GG, Gladman DD, Gómez-Reino J, Papp K, Baratelle A, Xu W, Mudivarthi S, Mack M, Rahman MU, Xu Z, Zrubek J, Beutler A. Golimumab in psoriatic arthritis: One-year clinical efficacy, radiographic, and safety results from a phase III, randomized, placebo-controlled trial.

Arthritis Rheum 2012;64:2504-17

Hermann K-G, Baraliakos X, [van der Heijde DMFM](#), Jurik A-G, Landewé R, Marzo-Ortega H, Østergaard M, Rudwaleit M, Sieper J, Braun J. Descriptions of spinal MRI lesions and definition of a positive MRI of the spine in axial spondyloarthritis: A consensual approach by the ASAS/OMERACT MRI study group.

Ann Rheum Dis 2012;71:1278-88 Chung H-Y, Machado P, [van der Heijde D](#), D'Agostino M-A, Dougados M. Smokers in early axial spondyloarthritis have earlier disease onset, more disease activity, inflammation and damage, and poorer function and health-related quality of life: results from the DESIR cohort.

Ann Rheum Dis published 2012;71:809-16

Haugen IK, Bøyesen P, Slatkowsky-Christensen B, Sesseng S, [van der Heijde D](#), Kvien TK. Associations between MRI-defined synovitis, bone marrow lesions and structural features and measures of pain and physical function in hand osteoarthritis.

Ann Rheum Dis 2012;71:899-904

Braun J, Baraliakos X, Hermann K-G, [van der Heijde D](#), Inman RD, Deodhar AA, Baratelle A, Xu S, Xu W, Hsu B. Golimumab reduces spinal inflammation in ankylosing spondylitis: MRI results of the randomised, placebo- controlled GO-RAISE study.

Ann Rheum Dis 2012;71:878-84

[van der Heijde D](#). Remission by imaging in rheumatoid arthritis: should this be

the ultimate goal?

Ann Rheum Dis 2012;71:i89-i92

Mandl P, Balint PV, Brault Y, Backhaus M, D'Agostino M-A, Grassi W, [van der Heijde D](#), de Miguel D, Wakefield RJ, Logeart I, Dougados M. Metrologic properties of ultrasound versus clinical evaluation of synovitis in rheumatoid arthritis: Results of a multicenter, randomized study.

Arthritis Rheum 2012;64:1272-82

van Tubergen A, Ramiro S, [van der Heijde D](#), Dougados M, Mielants H, Landewé R. Development of new syndesmophytes and bridges in ankylosing spondylitis and their predictors: a longitudinal study.

Ann Rheum Dis 2012;71:518-23 Lillegraven S, [van der Heijde D](#), Uhlig T, Kvien TK, Haavardsholm E. What is the clinical relevance of erosions and joint space narrowing in RA?

Nat Rev Rheumatol 2012;8:117-20

[van der Heijde D](#), Machado P, Braun J, Hermann K-GA, Baraliakos X, Hsu B, Baker D, Landewé R. MRI inflammation at the vertebral unit only marginally predicts new syndesmophyte formation: a multilevel analysis in patients with ankylosing spondylitis.

Ann Rheum Dis 2012;71:369-73

Haugen IK, Bøyesen P, Slatkowsky-Christensen B, Sesseng S, Bijsterbosch J, [van der Heijde D](#), Kvien TK. Comparison of features by MRI and radiographs of the interphalangeal finger joints in patients with hand osteoarthritis.

Ann Rheum Dis 2012;71:345-50

2011

Landewé R, [van der Heijde D](#). New considerations in the design of clinical trials for ankylosing spondylitis.

Clinical Investigation 2011;1:1505-12

Baraliakos X, Landewé R, [van der Heijde D](#), Listing J, Baker D, Hsu B, Braun J. The relationship of biomarkers and radiographic progression in patients treated with infliximab.

Eur Musc Rev 2011;6:94-95

Emery P, [van der Heijde D](#), Østergaard M, Conaghan PG, Genovese MC, Keystone EC, Fleischmann R, Hsia EC, Xu W, Xu S, Rahman MU. Exploratory analyses of the association of MRI with clinical, laboratory and radiographic findings in patients with rheumatoid arthritis.

Ann Rheum Dis 2011;70:2126-30

van Onna M, Jurik AG, [van der Heijde D](#), van Tubergen A, Heuft-Dorenbosch L, Landewé R. HLA-B27 and gender independently determine the likelihood of a positive MRI of the sacroiliac joints in patients with early inflammatory back pain: a 2-year MRI follow-up study.

Ann Rheum Dis 2011;70:1981-5

Chung HY, Machado P, [van der Heijde D](#), D'Agostino M-A, Dougados M. HLA-B27 positive patients differ from HLA-B27 negative patients in clinical presentation and imaging: results from the DESIR cohort of patients with recent onset axial

spondyloarthritis.

Ann Rheum Dis 2011;70:1930-6

Klarenbeek NB, Koevoets R, van der Heijde DMFM, Gerards AH, tenWolde S, Kerstens PJSM, Huizinga TWJ, Dijkmans BAC, Allaart CF. Association with joint damage and physical functioning of nine composite indices and the 2011 ACR/EULAR remission criteria in rheumatoid arthritis.

Ann Rheum Dis 2011;70:1815-21

Strand V, Kingsbury SR, Woodworth T, Landewé R, Ostergaard M, Peterfy C, van der Heijde D, D'Agostino MA, Maksymowych W, Tak PP, Wells G, Conaghan PG. OMERACT 10 Sharp symposium: Important findings in examination of imaging methods for measurement of joint damage in rheumatoid arthritis.

J Rheumatol 2011;38:2009-13

Landewé RB, Strand V, Conaghan PG, van der Heijde D. Damage and progression on radiographs in individual joints: data from pivotal randomized controlled trials.

J Rheumatol 2011;38:2018-22

Baraliakos X, van der Heijde D, Braun J, Landewé RB. OMERACT Magnetic Resonance Imaging initiative on structural and inflammatory lesions in ankylosing spondylitis – Report of a Special Interest Group at OMERACT 10 on sacroiliac joint and spine lesions.

J Rheumatol 2011;38:2051-4

Giles JT, van der Heijde DM, Bathon JM. Association of circulating adiponectin levels with progression of radiographic joint destruction in rheumatoid arthritis.

Ann Rheum Dis 2011;70 1562-1568

Lukas C, Combe B, Ravaud P, Sibilia J, Landewé R, D van der Heijde. Favorable effect of very early disease-modifying antirheumatic drug treatment on radiographic progression in early inflammatory arthritis data from the Etude et Suivi des Polyarthrites Indifférenciées Récentes (Study and Followup of Early Undifferentiated Polyarthritis).

Arthritis Rheum 2011;63:1804-11

Bijsterbosch J, Haugen IK, Malines C, Maheu E, Rosendaal FR, Watt I, Berenbaum F, Kvien TK, van der Heijde DM, Huizinga TWJ, Kloppenburg M. Reliability, sensitivity to change and feasibility of three radiographic scoring methods for hand osteoarthritis.

Ann Rheum Dis 2011;70 1465-7

van der Heijde D, Landewé R. Dramatic repair of joint damage in psoriatic arthritis.

J Rheumatol 2011;38:969-70

Emery P, Fleischmann R, van der Heijde D, Keystone EC, Genovese MC, Conaghan PG, Hsia EC, Xu W, Baratelle A, Beutler A, Rahman MU. The effects of golimumab on radiographic progression in rheumatoid arthritis: Results of randomized controlled studies of golimumab before methotrexate therapy and golimumab after methotrexate therapy.

Arthritis Rheum 2011;63:1200-10

Bøyesen P, Haavardsholm EA, Østergaard M, [van der Heijde D](#), Sesseng S, Kvien TK. MRI in early rheumatoid arthritis: synovitis and bone marrow oedema are independent predictors of subsequent radiographic progression. *Ann Rheum Dis* 2011;70:428-433

Bøyesen P, Haavardsholm EA, [van der Heijde D](#), Østergaard M, Berner Hammer H, Sesseng S, Kvien TK. Prediction of MRI erosive progression: a comparison of modern imaging modalities in early rheumatoid arthritis patients. *Ann Rheum Dis* 2011;70:176-9

Grandaunet B, Syversen SW, Hoff M, Sundan A, Haugeberg G, [van der Heijde D](#), Kvien TK, Standal T. Association between high plasma levels of hepatocyte growth factor and progression of radiographic damage in the joints of patients with rheumatoid arthritis. *Arthritis Rheum* 2011;63:662-9

[van der Heijde D](#). Erosions versus joint space narrowing in rheumatoid arthritis: what do we know? *Ann Rheum Dis* 2011;70:i116-i118

Hashimoto J, Garnero P, [van der Heijde D](#), Miyasaka N, Yamamoto K, Kawai S, Takeuchi T, Yoshikawa H, Nishimoto N. Humanized anti-interleukin-6-receptor antibody (tocilizumab) monotherapy is more effective in slowing radiographic progression in patients with rheumatoid arthritis at high baseline risk for structural damage evaluated with levels of biomarkers, radiography, and BMI: data from the SAMURAI study. *Mod Rheumatol*. 2011;21:10-5

Koevoets R, Machado P, Bombardier C, [van der Heijde DM](#). The value of conventional radiographs in undifferentiated arthritis: a systematic review. *J Rheumatol* 2011;38:Suppl 87:26-30

Machado PMMS, Koevoets R, Bombardier C, [van der Heijde DM](#). The value of magnetic resonance imaging and ultrasound in undifferentiated arthritis: a systematic review. *J Rheumatol* 2011;38:Suppl 87:31-7

Landewé R, [van der Heijde D](#). Joint space narrowing, cartilage and physical function: are we deceived by measurements and distributions? *Ann Rheum Dis* 2011;70:717-18

Haugen IK, Lillegraven S, Slatkowsky-Christensen B, Haavardsholm EA, Sesseng S, Kvien TK, [van der Heijde D](#), Boyesen P. Hand osteoarthritis and MRI: development and first validation step of the proposed Oslo Hand Osteoarthritis MRI score. *Ann Rheum Dis* 2011;70:1033-8

2010

Syversen SW, Goll GL, [van der Heijde D](#), Landewé R, Lie BA, Ødegard S, Uhlig T, Gaarder PI, Kvien TK. Prediction of radiographic progression in rheumatoid arthritis and the role of antibodies against mutated citrullinated vimentin: results from a 10-year prospective study. *Ann Rheum Dis* 2010;69:345-351

Emery P, Breedveld F, [van der Heijde D](#), Ferraccioli G, Dougados M, Robertson D, Pedersen R, Koenig AS, Freundlich B, for the Combination of Methotrexate and Etanercept in Early Rheumatoid Arthritis Trial Group. Two-year clinical and radiographic results with combination etanercept-methotrexate therapy versus monotherapy in early rheumatoid arthritis.

Arthritis Rheum 2010;62:674-82

van der Linden MPM, Boja R, Klarenbeek NB, Huizinga TWJ, [van der Heijde DM](#), van der Helm-van Mil AHM. Repair of joint erosions in rheumatoid arthritis: prevalence and patient characteristics in a large inception cohort.

Ann Rheum Dis 2010;69:727-9

Bridges Jr SL, Causey ZL, Burgos PI, Huynh BQN, Hughes LB, Danila M, van Everdingen A, Ledbetter S, Conn DL, Tamhane A, Westfall AO, Jonas BL, Callahan LF, Smith EA, Brasington R, Moreland LW, Alarcón GS, [van der Heijde DM](#). Radiographic severity of Rheumatoid Arthritis in African Americans: Results from a multicenter observational study.

Arthritis Care & Research 2010;62:624-31

Lukas C, [van der Heijde D](#), Fatenejad S, Landewé R. Repair of erosions occurs almost exclusively in damaged joints without swelling.

Ann Rheum Dis 2010;69:851-5 Syversen SW, Haavardsholm EA, Boyesen P, Goll GL, Okkenhaug C, Gaarder PI, [van der Heijde D](#), Kvien TK. Biomarkers in early rheumatoid arthritis: longitudinal associations with inflammation and joint destruction measured by magnetic resonance imaging and conventional radiographs.

Ann Rheum Dis 2010;69:845-50 van der Woude D, Syversen SW, van der Voort EIH, Verpoort KN, Goll GL, van der Linden MPM, van der Helm-van Mil AHM, [van der Heijde DMFM](#), Huizinga TWJ, Kvien TK, REM Toes. The ACPA isotype profile reflects long-term radiographic progression in rheumatoid arthritis.

Ann Rheum Dis 2010;69:1110-6 Machado P, Landewé R, Braun J, Hermann KG, Baker D, [van der Heijde D](#). Both structural damage and inflammation of the spine contribute to impairment of spinal mobility in patients with ankylosing spondylitis.

Ann Rheum Dis 2010;69:1465-70

[van der Heijde D](#), Breedveld FC, Kavanaugh A, Keystone EC, Landewé R, Patra K, Pangan AL. Disease Activity, Physical Function, and Radiographic Progression After Longterm Therapy with Adalimumab Plus Methotrexate: 5-Year Results of PREMIER.

J Rheumatol 2010; Oct 1 Epub

Zhang J, Redden DT, McGwin G Jr, Callahan LF, Smith EA, Alarcón GS, Moreland LW, [van der Heijde DM](#), Brown EE, Arnett DK, Mikuls TR, Bridges SL Jr, for the Consortium for the Longitudinal Evaluations of African Americans with Early Rheumatoid Arthritis Investigators. Generalized bone loss as a predictor of three-year radiographic damage in African American patients with recent-onset rheumatoid arthritis.

Arthritis Rheum 2010;62:2219-26 van der Helm-van Mil AHM, Knevel R, [van der Heijde D](#), Huizinga TWJ. How to avoid phenotypic misclassification in using joint destruction as an outcome measure for rheumatoid arthritis

Rheumatology 2010;49:1429-35

Dougados M, Jousse-Joulin S, Mistretta F, D'Agostino M-A, Backhaus M, Bentin J, Chales G, Chary-Valckenaere I, Conaghan P, Etchepare F, Gaudin P, Grassi W, [van der Heijde D](#), Sellam J, Naredo E, Szkudlarek M, Wakefield R, Saraux A. Evaluation

of several ultrasonography scoring systems of synovitis and comparison to clinical examination: Results from a prospective multi-center study of rheumatoid arthritis.

Ann Rheum Dis 2010;69:828-35

van der Heijde D, Rudwaleit M, Landewe RBM, Sieper J. Justification for including MRI as a tool in the diagnosis of axial SpA.

Nat Rev Rheumatol 2010;6:670–2 (doi:10.1038/nrrheum.2010.160)

Klarenbeek NB, Guler-Yuksel M, van der Heijde DMFM, Hulsmans HMJ, Kerstens PJSM, Molenaar THE, de Sonnaville PBJ, Huizinga TWJ, Dijkmans BAC, Allaart CF. Clinical synovitis in a particular joint is associated with progression of erosions and joint space narrowing in that same joint, but not in patients initially treated with infliximab.

Ann Rheum Dis published 2010;69 2107-13

2009

Vosse D, Landewé R, van der Heijde D, van der Linden S, van Staa T-P, Geusens P. Ankylosing spondylitis and the risk of fracture: results from a large primary care-based nested case control study.

Ann Rheum Dis 2009;68:1839-42

Landewé R, van der Heijde D. A systematic comparison of rheumatoid arthritis and ankylosing spondylitis: Structural outcomes.

Clin Exp Rheumatol 2009;27 (4 suppl 55):S102-S107 Syversen SW, Landewé R, van der Heijde D, Bathon JM, Boers M, Bykerk VP, Fitzgerald O, Gladman DD, Garnero P, Geusens P, Gladman DD, El-Gabalawy H, Inman RD, Kraus V, Kvien TK, Mease PJ, Østergaard M, Ritchlin CJ, Tak P-P, Taylor WJ, Maksymowych WP. Testing of the OMERACT 8 Draft Validation Criteria for a Soluble Biomarker Reflecting Structural Damage in RA: A Systematic Literature Search on 5 Candidate Biomarkers.

J Rheumatol 2009;36:1769-84

Maksymowych WP, Landewé R, Tak P-P, Ritchlin CJ, Østergaard M, Mease PJ, El-Gabalawy H, Garnero P, Gladman DD, Fitzgerald O, Aletaha D, Bykerk VP, Bathon JM, Syversen SW, Boers M, Geusens P, Inman RD, Kraus V, Kvien TK, Taylor WJ, Wells GA, van der Heijde D. Reappraisal of OMERACT 8 Draft Validation Criteria for a Soluble Biomarker Reflecting Structural Damage Endpoints in RA, PsA, and Spondyloarthritis: The OMERACT 9 v2 Criteria.

J Rheumatol 2009;36:1785-91 Maksymowych WP, Fitzgerald O, Wells GA, Gladman DD, Landewé R, Østergaard M, Taylor WJ, Christensen R, Tak P-P, Boers M, Syversen SW, Bathon JM, Ritchlin CJ, Mease PJ, Bykerk VP, Garnero P, Geusens P, El-Gabalawy H, Aletaha D, Inman RD, Byers Kraus V, Kvien TK, van der Heijde D. Proposal for Levels of Evidence Schema for Validation of a Soluble Biomarker Reflecting Damage Endpoints in RA, PsA, and AS, and Recommendations for Study Design.

J Rheumatol 2009;36:1792-9 Thabet M, Huizing TWJ, van der Heijde DM, van der Helm-van Mil AHM. The prognostic value of baseline erosions in undifferentiated arthritis.

Arthr Res Ther 2009;11:R155

van der Heijde D, Burmester GR, Melo-Gomes JR, Codreanu C, Martin Mola E, Pedersen R, Robertson D, Chang D, Koenig AS, Freundlich B. Inhibition of radiographic progression with combination etanercept and methotrexate in patients with moderately active rheumatoid arthritis previously treated with monotherapy.

Ann Rheum Dis 2009;68:1113-8

Berner Hammer H, Ødegard S, Syversen SW, Landewé R, van der Heijde D, Uhlig T, Møwinckel, Kvien TK. Calprotectin (a major S100 leukocyte protein) predicts 10-year radiographic progression in patients with rheumatoid arthritis.

Ann Rheum Dis 2010;69:150-4

Lukas C, Landewé RBM, Fatenejad S, van der Heijde DMFM. Subtle changes in individual joints result in both positive and negative change scores in a patient: Results from a clinical trial in patients with rheumatoid arthritis.

Ann Rheum Dis 2009;68:1691-5 Haavardsholm A, Østergaard M, Berner Hammer H, Boyesen P, Boonen A, van der Heijde D, Kvien TK. Monitoring anti-TNF{alpha} treatment in RA: Responsiveness of magnetic resonance imaging and ultrasonography of the dominant wrist compared to conventional measures of disease activity and structural damage.

Ann Rheum Dis 2009;68:1572-9

Smolen JS, van der Heijde DMFM, Aletaha D, Xu S, Han J, Baker D, St.Clair EW. Progression of radiographic joint damage in rheumatoid arthritis: Independence of erosions and joint space narrowing.

Ann Rheum Dis 2009;68:1535-40

Hashimoto J, Garnero P, van der Heijde D, Miyasaka N, Yamamoto K, Kawai S, Takeuchi T, Yoshikawa H, Nishimoto N. A combination of biochemical markers of cartilage and bone turnover, radiographic damage and body mass index to predict the progression of joint destruction in patients with rheumatoid arthritis treated with disease-modifying anti-rheumatic drugs.

Mod Rheumatol 2009;19:273-82 Smolen JS, Landewé RBM, Mease PJ, Brzezicki J, Mason D, Luijckens K, van Vollenhoven RF, Kavanaugh A, Schiff MH, Burmester GR, Strand V, Vencovsky J, van der Heijde DMFM. Efficacy and Safety of Certolizumab Pegol Plus Methotrexate in Active Rheumatoid Arthritis: The RAPID 2 Study. A randomized controlled. Trial.

Ann Rheum Dis 2009;68 797-804

Halvorsen EH, Haavardsholm EA, Pollmann S, Boonen A, van der Heijde D, Kvien TK, Molberg Ø. Serum IgG antibodies to Peptidylarginine Deiminase 4 predict radiographic progression in rheumatoid arthritis patients treated with TNF-{alpha} blocking agents.

Ann Rheum Dis 2009;68:249-52

Landewé R, Dougados M, Mielants H, van der Tempel H, van der Heijde D. Physical function in ankylosing spondylitis is independently determined by both disease activity and radiographic damage of the spine.

Ann Rheum Dis 2009;68 863-867

Smolen J, Han C, van der Heijde D, Emery P, Bathon J, Keystone E, Maini R, Kalden J, Aletaha D, Baker D, Han J, Bala M, St.Clair E. Radiographic changes in rheumatoid arthritis patients attaining different disease activity states with methotrexate monotherapy and infliximab plus methotrexate: the impacts of remission and TNF-blockade.

Ann Rheum Dis 2009;68 823-827

Hafstrom I, Albertsson K, Boonen A, van der Heijde D, Landewé R.B.M, Svensson B, Remission achieved after 2 years treatment with low dose prednisolone in addition to DMARD's in early RA is associated with reduced joint destruction still present after 4 years: an open two year continuation study.

Ann Rheum Dis 2009;68:508-513

Hoff M, Haugeberg G, Ødegard S, Syversen SW, Landewé R, van der Heijde D, Kvien K. Cortical hand bone loss after one year in early rheumatoid arthritis predicts radiographic hand joint damage at 5 and 10 year follow-up.
Ann Rheum Dis 2009;68:324-9

Syversen SW, Goll GL, van der Heijde D, Landewé R, Gaarder PI, Odegard S, Haavardsholm EA, Kvien TK. Cartilage and bone biomarkers in rheumatoid arthritis: prediction of 10-year radiographic progression.
J Rheumatol 2009;36:2-7

Berner Hammer H, Ødegard S, Syversen SW, Landewé R, van der Heijde D, Uhlig T, Møwinckel, Kvien TK. Calprotectin (a major S100 leukocyte protein) predicts 10-year radiographic progression in patients with rheumatoid arthritis.
Ann Rheum Dis published 18 December 2008, 10.1136/ard.2008.103739

Lukas C, Landewé RBM, Fatenejad S, van der Heijde DMFM. Subtle changes in individual joints result in both positive and negative change scores in a patient: Results from a clinical trial in patients with rheumatoid arthritis.
Ann Rheum Dis published 24 November 2008, 10.1136/ard.2008.097816

Haavardsholm A, Østergaard M, Berner Hammer H, Boyesen P, Boonen A, van der Heijde D, Kvien TK. Monitoring anti-TNF{alpha} treatment in RA: Responsiveness of magnetic resonance imaging and ultrasonography of the dominant wrist compared to conventional measures of disease activity and structural damage.
Ann Rheum Dis published 19 November 2008, 10.1136/ard.2008.091801

Smolen JS, van der Heijde DMFM, Aletaha D, Xu S, Han J, Baker D, St.Clair EW. Progression of radiographic joint damage in rheumatoid arthritis: Independence of erosions and joint space narrowing.
Ann Rheum Dis published 28 October 2008, 10.1136/ard.2008.094128

2008

Keystone E, van der Heijde D, Mason D Jr, Landewé R, van Vollenhoven R, Combe B, Emery P, Strand V, Mease P, Desai C, Pavelka K. Certolizumab Pegol Plus Methotrexate Is Significantly More Effective Than Placebo Plus Methotrexate in Active Rheumatoid Arthritis. Findings of a Fifty-Two-Week, Phase III, Multicenter, Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study.
Arthritis Rheum. 2008;58:3319-29

van der Heijde D, Landewé R, Baraliakos X, Houben H, van Tubergen A, Williamson P, Xu W, Baker D, Goldstein N, Braun J, the Ankylosing Spondylitis Study for the Evaluation of Recombinant Infliximab Therapy Study Group. Radiographic findings following two years of infliximab therapy in patients with ankylosing spondylitis.
Arthritis Rheum 2008;58: 3063-70

Vosse D, Landewé R, Garnero P, van der Heijde D, van der Linden S, Geusens P. Association of markers of bone- and cartilagedegradation with radiological changes at baseline and after 2 years follow-up in patients with ankylosing spondylitis.

Rheumatology 2008;47(8):1219-22

Lukas C, Sharp JT, Angwin J, Boers M, Duryea J, Hall JR, Kauffman JA, Landewé R, Langs G, Bernelot Moens HJ, Peloschek P, van der Heijde D. Automated measurement of joint space width in

small joints of patients with rheumatoid arthritis.

J Rheumatol. 2008;35(7):1288-1293

Antoni CE, Kavanaugh A, van der Heijde D, Beutler A, Keenan G, Zhou B, Kirkham B, Tutuncu Z, Burmester GR, Schneider U, Furst DE, Molitor J, Keystone E, Gladman DD, Manger B, Wassenberg S, Weier R, Wallace DJ, Weisman MH, Kalden JR, Smolen JS. Two-year efficacy and safety of infliximab treatment in patients with active psoriatic arthritis: Findings of the Infliximab Multinational Psoriatic Arthritis Controlled Trial (IMPACT).

J Rheum 2008;35(5):869-76

Cohen S, Dore R, Lane N, Ory P, Peterfy C, Sharp J, van der Heijde D, Zhou L, Tsuji W, Newmark R. Denosumab treatment effects on structural damage, bone mineral density, and bone turnover in rheumatoid arthritis: a twelve-month, multicenter, randomized, double-blind, placebo-controlled, phase II clinical trial.

Arthritis Rheum 2008;58(5):1299-309

van der Heijde D, Landewé R, Einstein S, Ory P, Vosse D, Ni L, Lin S.-L, Tsuji W, Davis Jr. J. Radiographic progression of ankylosing spondylitis after up to two years of treatment with etanercept.

Arthritis Rheum 2008;58(5):1324-31

Halvorsen EH, Pollmann S, Gilboe IM, van der Heijde D, Landewé R, Ødegard S, Kvien TK, Molberg, O. Serum IgG antibodies to peptidylarginine deiminase 4 in rheumatoid arthritis and associations with disease severity.

Ann Rheum Dis 2008;67(3):414-7

Dias EM, Lukas C, Landewé R, Fatenejad S, van der Heijde D. Reliability and sensitivity to change of the Simple Erosion Narrowing Score compared with the Sharp-van der Heijde method for scoring radiographs in rheumatoid arthritis.

Ann Rheum Dis 2008;67:375-9

Mikuls TR, Hughes LB, Westfall AO, Holers VM, Parrish L, van der Heijde D, van Everdingen M, Alarcon GS, Conn DL, Jonas B, Callahan LF, Smith EA, Gilkeson G, Howerd G, Moreland LW, Bridges Jr SL. Cigarette smoking, disease severity, and autoantibody expression in African American with recent-onset rheumatoid arthritis.

Ann Rheum Dis 2008;67 1529-1534

van der Heijde D, Landewé R, van Vollenhoven R, Fatenejad S, Klareskog L. Level of radiographic damage and radiographic progression are determinants of physical function: a longitudinal analysis of the TEMPO trial.

Ann Rheum Dis 2008;67:1267-70

Syversen SW, Gaarder PI, Goll GL, Odegard S, Haavardsholm EA, Mowinckel P, van der Heijde D, Landewé R, Kvien TK. High anti-cyclic citrullinated peptide levels and an algorithm of four variables predict radiographic progression in patients with rheumatoid arthritis: results from a 10-year longitudinal study.

Ann Rheum Dis 2008;67:212-7 van der Heijde D, Burmester G, Melo-Gomes J, Codreanu C, Martin Mola E, Pedersen R, Freundlich B, Chang DJ. For the Etanercept Study 400 investigators. The safety and efficacy of adding etanercept to methotrexate or methotrexate to etanercept in moderately active rheumatoid arthritis patients previously treated with monotherapy.

Ann Rheum Dis 2008;67:182-8

Dias EM, Lukas C, Landewé R, Fatenejad S, van der Heijde D. Reliability and sensitivity to change of the Simple Erosion Narrowing Score compared with the Sharp-van der Heijde method for scoring radiographs in rheumatoid arthritis. *Ann Rheum Dis* 2008; 67(3):375-379

2007

Lukas C, Braun J, van der Heijde D, Hermann KG, Rudwaleit M, Ostergaard M, Oostveen A, O'Connor P, Maksymowych WP, Lambert RG, Jurik AG, Baraliakos X, Landewé R; ASAS/OMERACT MRI in AS Working Group. Scoring inflammatory activity of the spine by magnetic resonance imaging in ankylosing spondylitis: a multireader experiment. *J Rheumatol* 2007;34(4):862-70

van der Heijde D, Landewé R, Hermann KG, Rudwaleit M, Ostergaard M, Oostveen A, O'Connor P, Maksymowych WP, Lambert RG, Lukas C, Jurik AG, Boers M, Baraliakos X, Braun J; ASAS/OMERACT MRI in AS Working Group. Is there a preferred method for scoring activity of the spine by magnetic resonance imaging in ankylosing spondylitis? *J Rheumatol*. 2007 Apr;34(4):871-3

Sharp JT, Angwin J, Boers M, Duryea J, von Ingersleben G, Hall JR, Kauffman JA, Landewé R, Langs G, Lukas C, Maillefert JF, Bernelet Moens HJ, Peloschek P, Strand V, van der Heijde D. Computer based methods for measurement of joint space width: update of an ongoing OMERACT project. *J Rheumatol*. 2007 Apr; 34(4):874-83

Lie BA, Viken MK, Odegard S, van der Heijde D, Landewé R, Uhlig T, Kvien TK. Associations between the PTPN22 1858C>T polymorphism and radiographic joint destruction in patients with rheumatoid arthritis: Results from a 10-year longitudinal study. *Ann Rheum Dis*;66:1604-09

Nishimoto N, Hashimoto J, Miyasaka N, Yamamoto K, Kawai S, Takeuchi T, Murata N, van der Heijde D, Kishimoto T. Study of Active Controlled Monotherapy Used for Rheumatoid Arthritis, an IL-6 inhibitor (SAMURAI): - Evidence of clinical and radiographic benefit from an X-Ray Reader-Blinded Randomized Controlled Trial of Tocilizumab. *Ann Rheum Dis*. 2007;66:1162-7

van der Heijde D, Landewé R, Boonen A, Einstein SG, Herborn G, Rau R, Wassenberg S, Weissman BV, Winalski CS, Sharp JT. Expert agreement confirms that negative changes in hand and foot radiographs are a surrogate for repair in patients with rheumatoid arthritis. *Arthritis Res Ther*. 2007 Jul 2;9(4):R62

van der Heijde D, Kavanaugh A, Gladman DD, Antoni C, Krueger GG, Guzzo C, Zhou B, Dooley LT, de Vlam K, Geusens P, Birbara C, Halter D, Beutler A; IMPACT 2 Study Group Members of the IMPACT 2 Study Group are shown in Appendix A. Infliximab inhibits progression of radiographic damage in patients with active psoriatic arthritis through one year of treatment: Results from the induction and maintenance psoriatic arthritis clinical trial 2. *Arthritis Rheum*. 2007 Jul 30;56980:2698-2707

Van der Heijde D, Klareskog L, Landewé R, Bruyn GA, Cantagrel A, Durez P, Herrero-Beaumont G, Molad Y, Codreanu C, Valentini G, Zahora R, Pedersen R, MacPeck D, Wajdula S, Fatenejad S.; TEMPO Study Investigators. Disease remission and

sustained halting of radiographic progression with combination etanercept and methotrexate in patients with rheumatoid arthritis.

Arthritis Rheum. 2007;56:3883-6

2006

Heuft-Dorenbosch L, Weijers R, Landewé R, van der Linden S, van der Heijde D. Magnetic Resonance Imaging changes of sacroiliac joints in patients with recent-onset inflammatory back pain: inter-reader reliability and prevalence of abnormalities.

Arthritis Res Therapy 2006;8:R11

Ødegård S, Landewé R, van der Heijde D, Kvien T, Mowinckel P, Uhlig T. Association of early radiographic damage with impaired physical function in rheumatoid arthritis: a ten-year, longitudinal observational study in 238 patients.

Arthritis Rheum. 2006 Jan;54(1):68-75

Kavanaugh A, Antoni C, Gladman D, Wassenberg S, Zhou B, Beutler A, Keenan G, Burmester G, Furst D, Weisman M, Kalden J, Smolen J, van der Heijde D. The Infliximab Multinational Psoriatic Arthritis Controlled Trial (IMPACT): Results of radiographic analyses after 1 year.

Ann Rheum Dis published on 13 February 2006 as 10.1136/ard.2005.04565 Heuft-Dorenbosch L, Landewé R, Weijers R, Wanders A, Houben H, van der Linden S, van der Heijde D. Combining information obtained from MRI and conventional radiographs in order to detect sacroiliitis in patients with recent-onset inflammatory back pain.

Ann Rheum Dis 2006;65:804-808

Smolen JS, van der Heijde DM, St. Clair EW, Emery P, Bathon JM, Keystone E, Maini RN, Kalden JR, Schiff M, Baker D, Han C, Han J, Bala M for the active-controlled study of patients receiving infliximab for the treatment of rheumatoid arthritis of early onset (ASPIRE) study group. Predictors of joint damage in patients with early rheumatoid arthritis treated with high-dose methotrexate with or without concomitant infliximab: results from the ASPIRE trial.

Arthritis Rheum. 2006;54(3):702-10

van der Heijde D, Klareskog L, Rodriguez-Valverde V, Codreanu C, Bolosiu H, Melo-Gomes J, Tornero-Molina J, Wajdula J, Pedersen R and Fatenejad S for the TEMPO Study Investigators. Comparison of Etanercept and Methotrexate, Alone and Combined, in the Treatment of Rheumatoid Arthritis: Two-Year Clinical and Radiography Results from the TEMPO Study, a Double-Blind, Randomized Trial.

Arthritis Rheum 2006;54(4):1063-1074

Geusens P, Landewé R, Garnero P, Chen D, Dunstan D, Lems W, Stinissen P, van der Heijde D, van der Linden S, Boers. The ratio of circulating osteoprotegerin to RANKL in early rheumatoid arthritis predicts later joint destruction.

Arthritis Rheum 2006;54:1772-1777

Braun J, Landewé R, Hermann K-G, Han J, Yan S, Williamson P, van der Heijde D for the ASSERT Study Group. Major reduction in spinal inflammation in patients with ankylosing spondylitis after treatment with infliximab: Results of a multicenter, randomized, double-blind, placebo-controlled magnetic resonance imaging study.

Arthritis Rheum 2006;54(5):1646-1652

Mease P, van der Heijde D. Joint damage in psoriatic arthritis: how is it assessed and can it be prevented?
Int J Adv Rheumatol 2006;4(2):38-48

Landewé R, van der Heijde D, Klareskog L, van Vollenhoven R, Fatenejad S. Disconnect between inflammation and joint destruction after treatment with etanercept plus methotrexate: results from the trial of etanercept and methotrexate with radiographic and patient outcomes.
Arthritis Rheum 2006;54(10):3119-25

Heuft-Dorenbosch L, Weijers R, Landewé R, van der Linden S, van der Heijde D. Magnetic resonance imaging changes of sacroiliac joints in patients with recent-onset inflammatory back pain: inter-reader reliability and prevalence of abnormalities.
Arthritis Res Ther 2006;8(1):R11

van der Heijde D, Klareskog L, Rodriguez-Valverde V, Codreanu C, Bolosiu H, Melo-Gomes J, Tornero-Molina J, Wajdula J, Pedersen R, Fatenejad S; TEMPO Study Investigators. Comparison of etanercept and methotrexate, alone and combined, in the treatment of rheumatoid arthritis: two-year clinical and radiographic results from the TEMPO study, a double-blind, randomized trial.
Arthritis Rheum 2006;54(4):1063-74

2005

van der Heijde D, Sharp J, Wassenberg S, Gladman D. Psoriatic arthritis imaging: a review of scoring methods.
Ann Rheum Dis 2005;64 suppl 2:ii61-4

van der Heijde D, Landewé R, Klareskog L, Rodríguez-Valverde V, Settas L, Pedersen R, Fatenejad S.. Presentation and analysis of data on radiographic outcome in clinical trials: experience from the TEMPO study.
Arthritis Rheum 2005;52:49-60

Wanders A, van der Heijde D, Landewe R, Behier JM, Calin A, Olivieri I, Zeidler H, Dougados M. Nonsteroidal antiinflammatory drugs reduce radiographic progression in patients with ankylosing spondylitis: A randomized clinical trial.
Arthritis Rheum 2005;52:1756-1765

Bruynesteyn K, Boers M, Kostense P, van der Linden S, van der Heijde D. Deciding on progression of joint damage in paired films of individual patients: smallest detectable difference or change
Ann Rheum Dis 2005;64:179-182

Baraliakos X, Landewé R, Hermann K-G, Listing J, Golder W, Brandt J, Rudwaleit M, Bollow M, Sieper J, van der Heijde D, Braun J. Inflammation in ankylosing spondylitis – a systematic description of the extent and frequency of acute spinal changes using magnetic resonance imaging (MRI)
Ann Rheum Dis 2005;64:730-4

Baraliakos X, Hermann K-G, Landewé R, Listing J, Golder W, Brandt J, Rudwaleit M, Bollow M, Sieper J, van der Heijde D, Braun J. Assessment of acute spinal inflammation in patients with ankylosing spondylitis by magnetic resonance imaging (MRI): a comparison between contrast enhanced T1 and short-tau inversion recovery (STIR) sequences.

Ann Rheum Dis 2005;64:1141-1144

van der Heijde D, Landewe R. Imaging in spondylitis.

Curr Opin Rheumatol 2005;17(4):413-7

Smolen JS, Han C, Bala M, Maini RN, Kalden JR, van der Heijde D, Breedveld FC, Furst DE and Lipsky PE for the ATTRACT Study Group. Evidence of radiographic benefit of treatment with infliximab plus methotrexate in rheumatoid arthritis patients who had no clinical improvement: A detailed subanalysis of data from the anti-tumor necrosis factor trial in rheumatoid arthritis with concomitant therapy study
Arthritis Rheum 2005;52:1020-1030

van der Heijde D, Landewé R, van der Linden S. How should treatment effect on spinal radiographic progression in patients with ankylosing spondylitis be measured?

Arthritis Rheum 2005;52:1979-1985

Sharp JT, van der Heijde D, Angwin J, Duryea J, Bernelot Moens HJ, Jacobs JW, Maillfert JF, Strand CV. Measurement of Joint Space Width and Erosion Size

J Rheumatol 2005;32(12):2456-61

Landewé R, van der Heijde D. Presentation and analysis of radiographic data in clinical trials and observational studies.

Ann Rheum Dis 2005;64 suppl IV:i48-51

Svensson B, Boonen A, Albertsson K, van der Heijde D, Keller C, Hafström I for the BARFOT Study Group. Low-dose prednisolone in addition to the initial disease-modifying antirheumatic drug in patients with early active rheumatoid arthritis reduces joint destruction and increases the remission rate.

Arthritis Rheum 2005;52:3360-70

Landewé R, van der Heijde D. Radiographic progression in rheumatoid arthritis.

Clin Exp Rheum 2005;23(suppl 39):S63-S68

van der Heijde D, Landewé R, Hermann K-G, Jurik A-G, Maksymowych W, Rudwaleit M, O'Connor P, Braun J and the ASAS/OMERACT MRI in AS working group. Application of the OMERACT filter to scoring methods for MRI of the SI joints and the spine. Recommendations for a research agenda at OMERACT 7.

J Rheumatol 2005;32:2042-7

van der Heijde D, Landewé R, and the ASAS working group. Selection of a method for scoring radiographs for AS clinical trials by ASAS and OMERACT.

J. Reumatol 2005;32:2048-58

Landewé R, Hermann K-G, van der Heijde D, Baraliakos X, Jurik A-G, Lambert R, Østergaard M, Rudwaleit M, Salonen D, Braun J and the ASAS/OMERACT MRI in AS working group. Scoring sacro-iliac joints by magnetic resonance imaging, a multiple-reader reliability experiment.

J Rheumatol 2005;32:2050-5

2004

van der Heijde D. Overview of radiologic efficacy of new treatments.
Rheum Dis Clin North Am 2004;30(2):285-93

Bruynesteyn K, Landewé R, van der Linden S, van der Heijde D. Radiography as primary outcome in rheumatoid arthritis: acceptable sample sizes for trials with 3 months' follow-up.
Ann Rheum Dis 2004;63(11):1413-1418

Landewé R, van der Heijde D. Radiographic progression depicted by probability plots: presenting data with optimal use of individual values.
Arthritis Rheum 2004;50(3):699-706

Spoorenberg A, de Vlam K, van der Linden S, Dougados M, Mielants H, van de Tempel H, van der Heijde D. Radiological scoring methods in ankylosing spondylitis. Reliability and change over 1 and 2 years.
J Rheumatol 2004;31(1):125-32.

van der Heijde D. Assessment of radiographs in longitudinal observational studies.
J Rheum 2004;31 Suppl 69:46-47

van der Heijde D, Kalden JR, Scott D, Smolen J, Strand V. Long-term evaluation of radiographic disease progression in a subset of patients with rheumatoid arthritis treated with leflunomide beyond 2 years: concise report.
Ann Rheum Dis 2004;63(6):737-9

Bruynesteyn K, van der Linden S, Landewé R, Gubler F, Weijers R, van der Heijde D. Progression of RA on plain films judged differently by expert radiologists and rheumatologists.
J Rheumatol 2004;31(6):1088-94

Braun J, Baraliakos X, Golder W, Hermann K-G, Listing J, Brandt J, Rudwaleit M, Zühlsdorf S, Bollow M, Sieper J, van der Heijde D. Analysing chronic spinal changes in ankylosing spondylitis - a systematic comparison of conventional x-rays with magnetic resonance imaging (MRI) using established and new scoring systems.
Ann Rheum Dis 2004;63 :1046-1055

Maini RN, Breedveld FC, Kalden JR, Smolen JS, Furst D, Weisman MH, St Clair EW, Keenan GF, van der Heijde D, Marsters PA, Lipsky PE; Anti-Tumor Necrosis Factor Trial in Rheumatoid Arthritis with Concomitant Therapy Study Group. Sustained improvement over two years in physical function, structural damage, and signs and symptoms among patients with rheumatoid arthritis treated with infliximab and methotrexate.
Arthritis Rheum 2004;50(4):1051-65.

Wanders A, Landewé R, Spoorenberg A, Dougados M, van der Linden S, Mielants H, van de Tempel H, van der Heijde D. What is the most appropriate radiologic scoring method for Ankylosing Spondylitis? A comparison of the available methods based on the outcome measures in rheumatology clinical trials filter.
Arthritis Rheum 2004;50(8):2622-32

Welsing P, Landewé R, van Riel P, Boers M, van Gestel A, van der Linden S, Swinkels H, van der Heijde D. The relationship between disease activity and radiologic progression in patients with rheumatoid arthritis: a longitudinal analysis.

Arthritis Rheum 2004;50(7):2082-93

van der Heijde D. Quantification of radiological damage in inflammatory arthritis: rheumatoid arthritis, psoriatic arthritis and ankylosing spondylitis.

Best Pract Res Clin Rheumatol 2004;18(6):847-860

St Clair EW, van der Heijde DM, Smolen JS, Maini RN, Bathon JM, Emery P, Keystone E, Schiff M, Kalden JR, Wang B, Dewoody K, Weiss R, Baker D. Combination of infliximab and methotrexate therapy for early rheumatoid arthritis: A randomized, controlled trial.

Arthritis Rheum 2004;50(11):3432-3443

Wanders A, Landewé R, Spoorenberg A, de Vlam K, Mielants H, Dougados M, van der Linden S, van der Heijde D. Scoring of radiographic progression in randomised clinical trials in ankylosing spondylitis: a preference for paired reading order.

Ann Rheum Dis 2004;63(12):1601-4.

Bruynesteyn K, van der Heijde D, Boers M, van der Linden S, Lassere M, van der Vleuten C. The Sharp/van der Heijde method out-performed the Larsen/Scott method on the individual patient level in assessing radiographs in early rheumatoid arthritis.

J Clin Epidemiol 2004;57(5):502-12

2003

Landewé R, Boers M, van der Heijde D. How to interpret radiologic progression in randomised clinical trial.

Rheumatology;2003;42:2-5

Braun J, Baraliakos X, Golder W, Brandt J, Rudwaleit M, Listing J, Bollow M, Sieper J, van der Heijde D. MRI examinations of the spine in patients with ankylosing spondylitis (AS) before and after infliximab therapy after evaluation of a new scoring system.

Arthritis Rheum 2003;48(4):1126-1136

Sharp J, van der Heijde D, Boers M, Boonen A, Bruynesteyn K, Emery P, Genant H, Herborn G, Jurik A, Lassere M, McQueen F, Østergaard M, Peterfy C, Rau R, Strand V, Wassenberg S, Weissman B for the Subcommittee on Healing of Erosions of the OMERACT Imaging Committee. Repair of erosions in rheumatoid arthritis does occur. Results from two studies by the OMERACT subcommittee on healing of erosions.

J Rheumatol 2003;30:5:1102-7

van der Heijde D, Sharp J, Rau R, Strand V on behalf of subcommittee on healing of erosions of the OMERACT imaging committee. OMERACT Workshop: Repair of structural damage in rheumatoid arthritis.

J Rheumatol 2003;30(5):1108-9

Landewé RB, van der Heijde, D. Is radiographic progression a realistic outcome measure in clinical trials with early inflammatory arthritis?

Clin Exp Rheum 2003;21(suppl. 31):S37-S41

van der Heijde D, Landewé R. Imaging: Do erosions heal?
Ann Rheum Dis 2003;62(suppl.II):ii10-12

2002

van der Heijde D, Simon L, Smolen J, Strand V, Sharp J, Boers M, Breedveld F, Weissmann M, Weinblatt M, Rau R, Lipsky P. How to report radiographic data in randomized clinical trials in rheumatoid arthritis? Guidelines from a roundtable discussion.
Arthritis Rheum 2002;47:215-8

Landewé R, Boers M, Verhoeven A, Westhovens R, van de Laar M, Markusse H, van Denderen C, Westedt M, Peeters A, Dijkmans B, Jacobs P, Boonen A, van der Heijde D, van der Linden S. COBRA-combination therapy in patients with early rheumatoid arthritis: long-term structural benefits of a brief intervention.
Arthritis Rheum 2002;2:347-56

Bruynesteyn K, van der Heijde D, Boers M, Saudan A, Peloso P, Paulus H, Houben H, Griffiths B, Edmonds J, Bresnihan B, Boonen A, van der Linden S. Determination of the minimal clinically important difference in rheumatoid arthritis joint damage of the Sharp/van der Heijde and Larsen/Scott scoring methods by clinical experts and comparison with the smallest detectable difference.
Arthritis Rheum 2002;46:913-20

van der Heijde D. Structural damage in rheumatoid arthritis as visualized through radiographs.
Arthritis Research 2002; 4 (suppl 2): S29-S33

Boers M, van der Heijde D. Prevention or retardation of joint damage in rheumatoid arthritis. Issues of definition, evaluation and interpretation of plain radiographs.
Drugs 2002;62:1717-24

Bruynesteyn K, van der Heijde D, Boers M, Verhoeven A, Boonen A, van der Linden S. Contribution of progression of erosive damage in previously eroded joints in early rheumatoid arthritis trials: COBRA trial as an example.
Arthritis Rheum 2002;47:532-6

Strand V, Landewé R, van der Heijde D. Using estimated yearly progression rates to compare radiographic data across recent randomised controlled trials in rheumatoid arthritis.
Ann Rheum Dis 2002;61(suppl II):ii64-ii66

Bruynesteyn K, van der Heijde D, Boers M, Saudan A, Peloso P, Paulus H, Houben H, Griffiths B, Edmonds J, Bresnihan B, Boonen A, van der Linden S. Detecting radiological changes in rheumatoid arthritis that are considered important by clinical experts: Influence of reading with or without known sequence.
J Rheumatol 2002;29(11):2306-12

2001

Bruynesteyn K, van der Heijde D, Boers M, Lassere M, Boonen A, Edmonds J, Houben H, Paulus H, Peloso P, Saudan A, van

der Linden Sj. Minimal clinically important difference in radiological progression of joint damage over 1 year in RA: Preliminary results of a validation study with clinical experts.
J Rheumatol 2001;28:904-910

Lassere M, van der Heijde D, Johnson K, Bruynesteyn K, Molenaar E, Boonen A, Verhoeven A, Emery P, Boers M. Robustness and generalisability of smallest detectable difference in radiological progression.
J Rheumatol 2001;28:911-3

Lassere M, van der Heijde D, Johnson KR, Boers M, Edmonds J. Reliability of measures of disease activity and disease damage in RA: Implications for the smallest detectable difference, the minimum clinical important difference, and analysis of treatment effects in randomized controlled trials.
J Rheumatol 2001;28:892-903 van der Heijde D, Lassere M, Edmonds J, Kirwan J, Strand V, Boers M for the OMERACT imaging task force. Minimal Clinically Important Difference in plain films in RA: group discussions, conclusions and recommendations.
J Rheumatol 2001;28:914-5

van der Heijde D. Radiographic assessment of joint disease in rheumatoid arthritis. Joint damage occurs early in the disease course.
J Musculoskeletal Med 2001;5 (suppl):S15-S20

van der Heijde D. Radiographic progression in rheumatoid arthritis: does it reflect outcome? Does it reflect therapy?
Ann Rheum Dis 2001;60:iii47-iii50

van der Heijde D. Aspects of the radiological evaluation of rheumatoid arthritis.
Ann Med Interne 2001; 152: 518-21

2000

van der Heijde D. How to read radiographs according to the Sharp|van der Heijde method.
J Rheumatol 2000;27:261-3 van der Heijde DMFM. Radiographic imaging: The "gold standard" for assessment of disease progression in rheumatoid arthritis.
Rheumatology 2000;39(suppl 1):9-16 Proudman SM, Conaghan PG, Richardson C, Green MJ, McGonagle D, Wakefield RJ, Griffiths B, Reece R, Miles S, Veale D, Martin M, Huston G, Gough A, Adebajo A, Helliwell P, van der Heijde D, Emery P. Treatment of poor-prognosis early rheumatoid arthritis: A randomized study of treatment with methotrexate, cyclosporin A and intraarticular corticosteroids compared with sulfasalazine alone.
Arthritis and Rheum 2000;43:1809-19

Hulsmans HMJ, Jacobs JWG, van der Heijde D, van Albada-Kuipers GA, Schenk Y, Bijlsma JWJ. Course of radiological damage during the first 6 years of rheumatoid arthritis.
Arthritis Rheum 2000;43:1927-40 Lipsky P, van der Heijde D, St. Clair EW, Furst DE, Breedveld F, Kalden JR, Smolen J, Weisman M, Emery P, Feldmann M, Harriman GR, Maini R for the Anti-Tumor Necrosis Factor Trial in Rheumatoid Arthritis with Concomitant Therapy Study Group. Infliximab and methotrexate in the treatment of rheumatoid arthritis.
New Eng J Med 2000;343:1594-1602

1999

van der Heijde D, Boers M, Lassere M. Methodological issues in radiographic scoring methods in RA.
J Rheumatol 1999;26:726-30

van der Heijde D. How to read radiographs according to the Sharp/van der Heijde method.
J Rheumatol 1999;743-5

Molenaar E, Edmonds J, Saudan A, Lassere M, Boers M, van der Heijde D. A practical exercise in reading RA radiographs by the Larsen and Sharp methods.
J Rheumatol 1999;26:746-8

van der Heijde D, Boonen A, Boers M, Kostense P, van der Linden S. Reading radiographs in chronological order, in pairs or as single films has important implications for the discriminative power of rheumatoid arthritis clinical trials.
Rheumatology 1999;38:1213-20

van der Heijde D, Dankert T, Nieman F, Rau R, Boers M. Reliability and sensitivity to change of a simplification of the Sharp/van der Heijde radiological assessment in rheumatoid arthritis.
Rheumatology 1999;38:941-7

1997

Boers M, Verhoeven A, Markusse H, van de Laar M, Westhovens R, van Denderen C, van Zeben J, Dijkmans B, Peeters A, Jacobs P, van den Brink H, Schouten H, van der Heijde D, Boonen A, van der Linden S. Combination of step-down corticosteroids, methotrexate and sulfasalazine, compared to sulfasalazine alone in early, active rheumatoid arthritis produces excellent 1-year disease control and low toxicity; the COBRA trial.
Lancet 1997;350:309-18

1996

van der Heijde DMFM. Plain X-rays in rheumatoid arthritis: overview of scoring methods, their reliability and applicability. In: Bailliere Clin Rheum. Modern imaging techniques. Part I: Traditional Methods. Bird HA and Dougados M [eds.].
Bailliere Tindall, London, 1996;10:435-53

1995

van der Heijde DMFM, van Leeuwen MA, van Riel PLCM, van de Putte LBA. Radiographic progression on X-rays of hands and feet during the first three years of rheumatoid arthritis measured according to Sharp's method [van der Heijde modification].
J Rheumatol 1995;22:1792-6

van der Heijde DMFM. Joint erosions and patients with early rheumatoid arthritis.
Br J Rheumatol 1995;34[suppl]:74-8

1992

van der Heijde DMFM, van Leeuwen MA, van Riel PLCM, Koster AM, van 't Hof MA, van Rijswijk MH, van de Putte LBA.

Biannual radiographic assessments of hands and feet in a three-year prospective followup of patients with early rheumatoid arthritis.

Arthritis Rheum 1992;35:26-34

1989

van der Heijde DMFM, van Riel PLCM, Nuver-Zwart HH, Gribnau FWJ, van de Putte LBA. Effects of hydroxychloroquine and sulfasalazine on progression of joint damage in rheumatoid arthritis.

Lancet 1989;I:1036-8

3. Chapters in books in relation to imaging

2008

van der Heijde D.

Outcome assessment of rheumatoid arthritis. Radiography.

In: Rheumatoid Arthritis. Hochberg MC, Silman AJ, Smolen JS, Weinblatt ME, Weisman MH (editors) Mosby Elsevier, Philadelphia 2008;260-6

Baratelle A, van der Heijde D.

Radiographic imaging end points in rheumatoid arthritis trials.

In: Clinical trials in rheumatoid arthritis and osteoarthritis. D. Reid, C. Miller (editors) Springer-Verlag London 2008:201-22

2005

van der Heijde D.

Radiological measurements of periarticular bone loss, erosions, and joint space: x-ray measurement of erosions and joint space narrowing.

In: Bone Disease in Rheumatology. M. Maricic, O. Gluck (editors), Lippincott, Williams & Wilkins, Philadelphia, 2005:53-58

2004

van der Heijde D.

Measurement of Radiological Outcomes.

In: Rheumatoid Arthritis. St. Clair E. W, Pisetsky D, Haynes B.F. (editors), Lippincott Williams & Wilkins, Philadelphia, 2004, 90-97

1995

van der Heijde DMFM.

Radiographs as an outcome measure in rheumatoid arthritis.

Rheumatology in Europe 1995;24(suppl 2):127-8

1993

van de Putte LBA, van Riel PLCM, Boerbooms AMTh, van der Heijde DMFM, Jeurissen MEC. DMARDs: do they retard progression of radiological joint damage in rheumatoid arthritis? Eular Publishers, 1993

1992 van de Putte LBA, van der Heijde DMFM, van Riel PLCM.

Sulfasalazin-Behandlung und radiologische Gelenkschädigung bei Polyarthritits.

In: Wagenhäuser FJ, ed. Salazopyrin EN - Standortbestimmung und Ausblick. Eular Publishers, 1992